

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 November 2001 (15.11.2001)

PCT

(10) International Publication Number
WO 01/85714 A1

(51) International Patent Classification⁷: **C07D 333/76**,
A61K 31/381, 31/453, 31/454, 31/4436, 31/4439, A61P
9/00, 9/12, 3/04, C07D 409/12, 413/12, 521/00, 417/12,
409/14

FOOTE, Kevin, Michael [GB/GB]; Alderley Park, Mac-
clesfield, Cheshire SK10 4TG (GB). **BRITTAIN, David,**
Robert [GB/GB]; Alderley Park, Macclesfield, Cheshire
SK10 4TG (GB).

(21) International Application Number: PCT/GB01/01899

(74) Agent: **BRYANT, Tracey**; AstraZeneca, Global Intellec-
tual Property, Alderley Park, Mereside, P.O. Box 272, Mac-
clesfield,, Cheshire SK10 4GR (GB).

(22) International Filing Date: 1 May 2001 (01.05.2001)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:
0010757.3 5 May 2000 (05.05.2000) GB

(71) Applicant (*for all designated States except MG, US*): **AS-
TRAZENECA AB** [SE/SE]; S-151 85 Sodertalje (SE).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for MG only*): **ASTRAZENECA UK LIM-
ITED** [GB/GB]; 15 Stanhope Gate, London W1Y 6LN
(GB).

(72) Inventors; and

Published:


— with international search report

(75) Inventors/Applicants (*for US only*): **BLOCK, Michael,**
Howard [GB/GB]; Alderley Park, Macclesfield, Cheshire
SK10 4TG (GB). **DONALD, Craig, Samuel** [GB/GB];
Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: AMINO SUBSTITUTED DIBENZOTHIOPHENE DERIVATIVES FOR THE TREATMENT OF DISORDERS ME-
DIATED BY THE NP Y5 RECEPTOR

WO 01/85714 A1



(57) Abstract: Compounds of formula (I),
wherein X is a group of formula (A) or (B) and
R¹, R², R³, R⁴, n, x, y and z are defined within
are described. Processes for their preparation
and their use in the treatment of disorders
mediated by the neuropeptide Y5 receptor in
a warm-blooded animal, such as a human being
are also described.

- 1 -

AMINO SUBSTITUTED DIBENZOTHIOPHENE DERIVATIVES FOR THE TREATMENT OF DISORDERS
MEDIATED BY THE NP Y5 RECEPTOR

This invention relates to compounds which antagonise the interaction between neuropeptide Y (NPY) and the neuropeptide Y5 (NPY-5) receptor sub-type. This invention
5 also relates to processes for the manufacture of NPY-5 receptor antagonists or agonists, pharmaceutically acceptable salts thereof, and to novel pharmaceutical compositions of NPY-5 receptor antagonists or agonists.

NPY is a 36 amino acid polypeptide which is a member of the pancreatic polypeptide family of regulatory peptides with widespread distribution throughout the mammalian system.
10 NPY is the most abundant neuropeptide in the central and peripheral nervous systems and has been shown to have powerful and complex effects on feeding, anxiety, circadian rhythms, reproduction, pituitary-adrenocortical axis function, memory retention, seizures, thermo-regulation, and cardiovascular and gastrointestinal functions. NPY interacts with a heterogeneous population of at least six receptor subtypes, Y₁-Y₆ which activate adenylate
15 cyclase via a G-protein. For reviews of NPY see: CRC Critical Reviews in Neurobiology. (1988) 4, 97-135; Regulatory Peptides (1996) 62, 1-11.

One of the most striking actions of NPY is induction of feeding in a variety of vertebrate species. Direct injection of NPY into the hypothalamus of satiated rats can increase food intake up to 10-fold over a 4 hour period and NPY is the only known peptide which can
20 cause animals to eat until they are obese. Recent studies on NPY have focussed on the identification of the NPY receptor responsible for the regulation of feeding. The NPY-5 receptor has been identified as the receptor most closely matching a proposed appetite receptor. The functional role of this receptor was addressed by receptor blockade studies. Intra-cerebro-ventricular injection of NPY-5 receptor antisense oligodeoxynucleotides
25 prevented the increase in hypothalamic NPY levels during food deprivation and inhibited fasting-induced food intake in rats [Schaffhauser et al (1997) Diabetes 46, 1792 - 1798]. Thus the NPY-5 receptor is a potential pharmacological target in the modulation of feeding disorders such as obesity. For reviews on the association between NPY and feeding see: Zimanyi et al (1998) Current Pharm Des 4, 349-66; Heinrichs et al (1998) Vitamins and
30 Hormones 54, 51-66.


Obesity is a large and ever expanding problem in affluent societies, which has reached epidemic proportions. According to the US Institute of Medicine, 59% of Americans are

- 2 -

clinically obese or at least 20% above their ideal body weight. Obesity is associated with susceptibility to a number of other conditions e.g. non-insulin-dependent diabetes, hypertension, dyslipidaemia and coronary heart disease. These conditions lead to reduction in life expectancy and decreased quality of life. The overall financial burden of obesity is difficult to quantify but it has been estimated that in the US it may account for 6-8% of total healthcare expenditure.



Thus there is need for pharmaceutical agents which have efficacy in the treatment of eating disorders such as obesity, anorexia and bulimia. Modulation of NPY activity through antagonism at the NPY-5 receptor offers one potential target for pharmacological intervention in these conditions.

Accordingly, the present invention provides a compound of formula (I):



wherein:

X is a group of formula (A) or (B):



R^1 is cyano, halo, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl, C_{1-4} alkoxy, N -(C_{1-4} alkyl)amino or N,N -(C_{1-4} alkyl)₂amino;

R^2 is halo, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl or C_{1-4} alkoxy;

A is $-NR^a-$, $-O-$ or a direct bond; wherein R^a is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl; wherein R^a may be optionally substituted by one or more R^5 ;

R^3 is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl or C_{2-10} alkynyl wherein R^3 may be optionally substituted by one or more R^6 ; or R^3 is carbocyclyl or heterocyclyl wherein R^3 may be optionally substituted on carbon by one or more R^7 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R^8 ;

- 3 -

R^4 is halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-4} alkoxycarbonyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, C_{1-4} alkanoylamino, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, 5 N -(carbocyclyl)carbamoyl, N,N -(carbocyclyl)₂carbamoyl, N -(heterocyclyl)carbamoyl, N,N -(heterocyclyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0 to 2, C_{1-4} alkoxycarbonyl, N -(C_{1-4} alkyl)sulphamoyl, N,N -(C_{1-4} alkyl)₂sulphamoyl, C_{1-4} alkylsulphonylamino or (nitrogen-linked heterocyclic ring)carbonyl;

R^5 and R^6 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, 10 amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, C_{1-6} alkanoylamino, C_{2-6} alkenyloxy, carbonyl, C_{1-6} alkoxycarbonyl, heterocyclyl, heterocyclyloxy, heterocyclylamino, heterocyclyl- N -(C_{1-6} alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl, heterocyclylcarbonylamino, heterocyclyloxy, carbocyclyl, carbocyclyloxy, 15 carbocyclylamino, carbocyclyl- N -(C_{1-6} alkyl)amino, carbocyclylsulphonyl, carbocyclylcarbonyl, carbocyclylcarbonylamino, carbocyclyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkoxycarbonylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkylS(O)_a wherein a is 0-2, N -(C_{1-6} alkyl)sulphamoylamino, N,N -(C_{1-6} alkyl)₂sulphamoylamino, C_{1-6} alkylsulphonylamino, 20 (C_{1-6} alkyl)sulphonyl- N -(C_{1-6} alkyl)amino, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl)₂sulphamoyl; wherein R^5 and R^6 may be optionally substituted on carbon by one or more R^9 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be independently optionally substituted by R^{10} ;

R^7 and R^9 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, 25 trifluoromethyl, trifluoromethoxy, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, C_{1-4} alkoxycarbonylamino, C_{2-4} alkenyloxy, carbonyl, C_{1-4} alkanoyl, C_{1-4} alkanoylamino, C_{1-4} alkanoyloxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0-2, N -(C_{1-4} alkyl)sulphamoylamino, 30 N,N -(C_{1-4} alkyl)₂sulphamoylamino, (C_{1-4} alkyl)sulphonylamino, (C_{1-4} alkyl)sulphonyl- N -(C_{1-4} alkyl)amino, N -(C_{1-4} alkyl)sulphamoyl, N,N -(C_{1-4} alkyl)₂sulphamoyl, heterocyclyl, heterocyclyloxy, heterocyclylamino,

- 4 -

heterocyclyl-*N*-(C₁₋₄alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl, heterocyclylcarbonylamino, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylamino, carbocyclyl-*N*-(C₁₋₄alkyl)amino, carbocyclylsulphonyl, carbocyclylcarbonyl, carbocyclylcarbonylamino, and carbocyclyloxycarbonyl; wherein R⁷ and

5 R⁹ may be independently optionally substituted on carbon by one or more R¹¹;

R⁸ and R¹⁰ are independently selected from C₁₋₄alkyl, C₁₋₄alkanoyl, sulphonyl, C₁₋₄alkylsulphonyl, C₁₋₄alkoxycarbonyl, carbamoyl, *N*-(C₁₋₄alkyl)carbamoyl, *N,N*-(C₁₋₄alkyl)₂carbamoyl, *N*-(C₁₋₄alkyl)sulphonyl, *N,N*-(C₁₋₄alkyl)₂sulphonyl, heterocyclyl, heterocyclylcarbonyl, heterocyclyloxycarbonyl, heterocyclylsulphonyl, carbocyclyl, carbocyclylcarbonyl, carbocyclyloxycarbonyl and carbocyclylsulphonyl; wherein

R⁸ and R¹⁰ may be independently optionally substituted on carbon by one or more R¹²;

R¹¹ and R¹² are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphonyl, methyl, ethyl, vinyl, allyl, methoxy, ethoxy, vinyloxy, allyloxy, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, *N*-methylcarbamoyl, *N,N*-dimethylcarbamoyl, methylthio, methylsulphonyl, mesyl, *N*-methylsulphonyl, *N,N*-dimethylsulphonyl, heterocyclyl, heterocyclyloxy, heterocyclylcarbonyl, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylcarbonyl and carbocyclyloxycarbonyl;

x is 0 - 4; wherein the values of R¹ may be the same or different;
 20 y is 0 - 3; wherein the values of R² may be the same or different;
 z is 0 - 3; wherein the values of R⁴ may be the same or different; and
 n is 0 - 2;

or a pharmaceutically acceptable salt, prodrug or solvate thereof;


with the provisos when X is a group of formula (A):

- 25 1) when R³ is a nitrogen linked heterocyclyl, A is a direct bond;
 2) when x and y are 0 and n is 2, the group R³-A-C(O)-NH- is not 2-formamido, 2-acetamido, 3-acetamido, 2-propionamido, 3-[2-(fur-2-ylcarbonylmethyl)acetamido], 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido), 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido),
 30 2-(2-phthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido), 2-(2-amino-2-benzylacetamido), 3-(2-phthalimidoacetamido), 3-(3-phthalimidopropionamido),

- 5 -

- 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido),
 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido),
 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-benzylacetamido) or
 2-(3,3,3-trifluoro-2-hydroxy-2-methylpropionamido);
- 5 3) when x and y are 0 and n is 1, the group $R^3-A-C(O)-NH-$ is not 3-acetamido,
 3-(2-phthalimidoacetamido), 3-(3-phthalimidopropionamido),
 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido),
 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido),
 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-isobutylacetamido) or
- 10 3-(2-amino-2-benzylacetamido);
- 4) when x and y are 0 and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-benzamido, 2-acetamido,
 2-benzyloxycarbonylamino, 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido),
 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido),
 2-(2-phthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido),
- 15 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido),
 2-(2-amino-2-benzylacetamido), 3-(trifluoroacetamido), 3-benzyloxycarbonylamino or
 4-t-butyloxycarbonylamino;
- 5) when $(R^1)_x$ is 7-fluoro, y is 0 and n is 2, the group $R^3-A-C(O)-NH-$ is not 3-acetamido;
- 6) when x is 0, $(R^2)_y$ is 1-cyano and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido;
- 20 7) when x is 0, $(R^2)_y$ is 3-bromo and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido or
 2-benzamido;
- 8) when x is 0, $(R^2)_y$ is 1-bromo and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido or
 2-benzamido; and
- 9) when x is 0, $(R^2)_y$ is 1-chloro or 4-chloro and n is 0, the group $R^3-A-C(O)-NH-$ is not
- 25 2-acetamido.

The numbering system for the dibenzothiophene ring used in the present specification is as follows:



For the avoidance of doubt, where any C₁₋₆alkyl is optionally substituted, this also includes the possibility of optional substitution on other groups that contain a C₁₋₆alkyl group, for example a C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, C₁₋₆alkanoylamino, C₁₋₆alkoxycarbonyl, *N*-(C₁₋₆alkyl)amino, *N,N*-di-(C₁₋₆alkyl)amino, C₁₋₆alkoxycarbonylamino, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-di-(C₁₋₆alkyl)carbamoyl or a C₁₋₆alkylS(O)_a wherein a is 0-2 group. Similarly, where any heterocyclyl or carbocyclyl may be optionally substituted this also includes the possibility of optional substitution on other groups that contain a heterocyclyl or carbocyclyl group, for example heterocyclyloxy carbonyl and carbocyclyl carbonyl.

10 In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. For example, "C₁₋₁₀alkyl", "C₁₋₆alkyl" and "C₁₋₄alkyl" include propyl, isopropyl and *t*-butyl. However, references to individual alkyl groups such as 'propyl' are specific for the straight chained version only and references to individual branched chain alkyl
15 groups such as 'isopropyl' are specific for the branched chain version only. A similar convention applies to other radicals, for example "phenylC₁₋₆alkyl" includes phenylC₁₋₄alkyl, benzyl, 1-phenylethyl and 2-phenylethyl. The term "halo" refers to fluoro, chloro, bromo and iodo.

Where optional substituents are chosen from "one or more" groups it is to be
20 understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups.

A "heterocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 3-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH₂-
25 group can optionally be replaced by a -C(O)- or a ring sulphur atom may be optionally oxidised to form the S-oxides. Preferably a "heterocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 5 or 6 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH₂- group can optionally be replaced by a -C(O)- or a ring
30 sulphur atom may be optionally oxidised to form the S-oxides. Examples and suitable values of the term "heterocyclyl" are thiazolidinyl, pyrrolidinyl, pyrrolinyl, 1,1-dioxotetrahydrothienyl, 2-pyrrolidone, 2-oxazolidinone, 4-thiazolidone, morpholino,

tetrahydropyranyl, piperidyl, piperazinyl, thiomorpholino, 1,1-dioxothiomorpholino, homopiperazinyl, thienyl, isoxazolyl, imidazolyl, pyrrolyl, thiadiazolyl, isothiazolyl, triazolyl, pyranyl, indolyl, pyrimidyl, pyrazinyl, pyridazinyl, pyridyl, 4-pyridone, quinolyl, 1,3-benzodioxolyl and 1-isoquinolone. Preferably the term "heterocyclyl" refers to

5 1,1-dioxotetrahydrothienyl, 2-pyrrolidone, 2-oxazolidinone, morpholino, tetrahydropyranyl, piperidyl, piperazinyl, 1,1-dioxothiomorpholino, imidazolyl, triazolyl and pyridyl. More preferably the term "heterocyclyl" refers to tetrahydropyranyl, morpholino, 2-oxopyrrolidinyl, piperidinyl, pyrrolidinyl, azetidyl, tetrahydrofuryl, 1,4-dioxo-8-azaspiro[4.5]decanyl, 1,1-dioxotetrahydrothienyl, 1,2,3,4,-tetrahydro-9H-pyrido[3,4-b]indole,

10 decahydroquinolin-1-yl, 1,2,5,6-tetrahydropyridyl, piperazinyl, pyridyl, 1,2,4-triazolyl, thienyl, 2-oxooxazolidinyl, imidazolyl, 1,1-dioxothiomorpholino, 2-oxo-1,2-dihydropyridyl, benzimidazolyl, pyrazolyl, succinimido, 2,4-dioxothiazolidinyl, furyl, pyridazinyl, 1,4-dihydrooxazin-2-one, 2-oxo-2,3-dihydrobenzimidazolyl, 2,3-dihydro-2-oxobenzimidazolyl, thieno[2,3-d]pyrimidinyl or pyrimidinyl.

15 A "nitrogen-linked heterocyclic ring" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 3-12 atoms of which at least one atom is nitrogen, which is linked via a nitrogen atom, wherein a -CH₂- group can optionally be replaced by a -C(O)- or a ring sulphur atom may be optionally oxidised to form the S-oxides. Examples and suitable values of a "nitrogen-linked heterocyclic ring" are morpholino, pyrrolidin-1-yl, imidazol-1-yl,

20 1,1-dioxothiomorpholino and 2-oxopyrrolidin-1-yl.

A "carbocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic carbon ring that contains 3-12 atoms. Preferably "carbocyclyl" is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. Suitable values for "carbocyclyl" include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl,

25 tetralinyl or indanyl. Particularly "carbocyclyl" is cyclopropyl or phenyl. More particularly "carbocyclyl" refers to cyclopropyl, cyclopentyl, cyclohexyl, bicyclo[2,2,1]heptyl or phenyl.

An example of "C₁₋₆alkanoyloxy" and "C₁₋₄alkanoyloxy" is acetoxy. Examples of "C₁₋₆alkoxycarbonyl" and "C₁₋₄alkoxycarbonyl" include methoxycarbonyl, ethoxycarbonyl, *n*- and *t*-butoxycarbonyl. Examples of "C₁₋₆alkoxycarbonylamino" and

30 "C₁₋₄alkoxycarbonylamino" include methoxycarbonylamino, ethoxycarbonylamino, *n*- and *t*-butoxycarbonylamino. Examples of "C₁₋₆alkoxy" and "C₁₋₄alkoxy" include methoxy, ethoxy and propoxy. Examples of "C₁₋₆alkanoylamino" and "C₁₋₄alkanoylamino" include formamido,

- acetamido and propionylamino. Examples of " $C_{1-6}alkylS(O)_a$ wherein a is 0 to 2" and " $C_{1-4}alkylS(O)_a$ wherein a is 0 to 2" include methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl and ethylsulphonyl. Examples of " $C_{1-6}alkylsulphonyl$ " include mesyl and ethylsulphonyl. Examples of " $C_{1-6}alkanoyl$ " and " $C_{1-4}alkanoyl$ " include propionyl and acetyl. Examples of " $N-(C_{1-6}alkyl)amino$ " and " $N-(C_{1-4}alkyl)amino$ " include methylamino and ethylamino. Examples of " $N,N-(C_{1-6}alkyl)_2amino$ " and " $N,N-(C_{1-4}alkyl)_2amino$ " include di-*N*-methylamino, di-*N*-ethylamino and *N*-ethyl-*N*-methylamino. Examples of " $C_{2-10}alkenyl$ " and " $C_{2-4}alkenyl$ " are vinyl, allyl and 1-propenyl. Examples of " $C_{2-10}alkynyl$ " and " $C_{2-4}alkynyl$ " are ethynyl, 1-propynyl and 2-propynyl. Examples of
- 10 " $C_{2-6}alkenyloxycarbonyl$ " and " $C_{2-4}alkenyloxycarbonyl$ " are vinyloxycarbonyl, allyloxycarbonyl and 1-propenyloxycarbonyl. Examples of " $N-(C_{1-6}alkyl)sulphamoyl$ " and " $N-(C_{1-4}alkyl)sulphamoyl$ " are *N*-(methyl)sulphamoyl and *N*-(ethyl)sulphamoyl. Examples of " $N-(C_{1-6}alkyl)_2sulphamoyl$ " and " $N-(C_{1-4}alkyl)_2sulphamoyl$ " are *N,N*-(dimethyl)sulphamoyl and *N*-(methyl)-*N*-(ethyl)sulphamoyl. Examples of " $N-(C_{1-6}alkyl)carbamoyl$ " and
- 15 " $N-(C_{1-4}alkyl)carbamoyl$ " are methylaminocarbonyl and ethylaminocarbonyl. Examples of " $N,N-(C_{1-6}alkyl)_2carbamoyl$ " and " $N,N-(C_{1-4}alkyl)_2carbamoyl$ " are dimethylaminocarbonyl and methylethylaminocarbonyl. Examples of "heterocycloxy" are pyridyloxy and thiazolyloxy. Examples of "heterocyclylcarbonyl" are pyrimidylcarbonyl and morpholinocarbonyl. Examples of "heterocyclyloxycarbonyl" are pyrrolidinylloxycarbonyl and pyranilyloxycarbonyl.
- 20 Examples of "carbocycloxy" are phenoxy and cyclopropyloxy. Examples of "carbocyclylcarbonyl" are benzoyl and cyclohexylcarbonyl. Examples of "carbocyclyloxycarbonyl" are phenoxycarbonyl and indanyloxycarbonyl. Examples of " $N-(carbocyclyl)carbamoyl$ " are *N*-phenylcarbamoyl and *N*-cyclopropylcarbamoyl. Examples of " $N,N-(carbocyclyl)_2carbamoyl$ " are *N,N*-diphenylcarbamoyl and
- 25 *N*-cyclohexyl-*N*-cyclopropylcarbamoyl. Examples of " $N-(heterocyclyl)carbamoyl$ " are *N*-pyridylcarbamoyl and *N*-furylcarbamoyl. Examples of " $N,N-(heterocyclyl)_2carbamoyl$ " are *N,N*-dipyridylcarbamoyl and *N*-pyrimidinyl-*N*-pyranilylcarbamoyl. Examples of " $C_{1-6}alkylsulphonylamino$ " and " $C_{1-4}alkylsulphonylamino$ " are mesylamino and isopropylsulphonylamino. Examples of " $(C_{1-6}alkyl)sulphonyl-N-(C_{1-6}alkyl)amino$ " and
- 30 $(C_{1-4}alkyl)sulphonyl-N-(C_{1-4}alkyl)amino$ are mesyl-*N*-methylamino and ethylsulphonyl-*N*-isopropylamino. Examples of "(nitrogen-linked heterocyclic ring)carbonyl" are morpholinocarbonyl and piperazin-1-ylcarbonyl. Examples of "heterocyclylamino" are

pyridylamino and furylamino. Examples of "heterocyclyl-*N*-(C₁₋₆alkyl)amino" and "heterocyclyl-*N*-(C₁₋₄alkyl)amino" are pyrimidinyl-*N*-methylamino and thienyl-*N*-isopropylamino. Examples of "heterocyclylsulphonyl" are pyridylsulphonyl and morpholin sulphonyl. Examples of "heterocyclylcarbonylamino" are morpholinocarbonylamino and thienylcarbonylamino. Examples of "carbocyclylamino" are cyclopropylamino and anilino. Examples of "carbocyclyl-*N*-(C₁₋₆alkyl)amino" are cyclopropyl-*N*-methylamino and *N*-methylanilino. Examples of "carbocyclylsulphonyl" are phenylsulphonyl and cyclohexylsulphonyl. Examples of "carbocyclylcarbonylamino" are benzoylamino and cyclopentylcarbonylamino. Examples of "*N*-(C₁₋₆alkyl)sulphamoylamino" and "*N*-(C₁₋₄alkyl)sulphamoylamino" are *N*-(methyl)sulphamoylamino and *N*-(ethyl)sulphamoylamino. Examples of "*N*-(C₁₋₆alkyl)₂sulphamoylamino" and "*N*-(C₁₋₄alkyl)₂sulphamoylamino" are *N,N*-(dimethyl)sulphamoylamino and *N*-(methyl)-*N*-(ethyl)sulphamoylamino.

A suitable pharmaceutically-acceptable salt of a compound of formula (I) is, for example, an acid-addition salt of a compound of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In addition a suitable pharmaceutically-acceptable salt of a compound of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

The compounds of the formula (I) may be administered in the form of a prodrug which is broken down in the human or animal body to give a compound of the formula (I). Examples of prodrugs include *in vivo* hydrolysable esters of a compound of the formula (I).

Various forms of prodrugs are known in the art. For examples of such prodrug derivatives, see:

- a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);
- b) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen and H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard

p. 113-191 (1991);

- c) H. Bundgaard, *Advanced Drug Delivery Reviews*, **8**, 1-38 (1992);
- d) H. Bundgaard, *et al.*, *Journal of Pharmaceutical Sciences*, **77**, 285 (1988); and
- e) N. Kakeya, *et al.*, *Chem Pharm Bull*, **32**, 692 (1984).

5 An *in vivo* hydrolysable ester of a compound of the formula (I) containing a carboxy or a hydroxy group is, for example, a pharmaceutically-acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically-acceptable esters for carboxy include C₁₋₆alkoxymethyl esters for example methoxymethyl, C₁₋₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl
10 esters, C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters, for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

15 An *in vivo* hydrolysable ester of a compound of the formula (I) containing a hydroxy group includes inorganic esters such as phosphate esters (including phosphoramidic cyclic esters) and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group/s. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxy-methoxy. A
20 selection of *in vivo* hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and N-(dialkylaminoethyl)-N-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl. Examples of substituents on benzoyl include morpholino and piperazino linked from a ring nitrogen atom via a methylene group to
25 the 3- or 4- position of the benzoyl ring.


It is to be understood that, insofar as certain of the compounds of formula I defined above may exist in optically active or racemic forms by virtue of one or more asymmetric carbon atoms, the invention includes in its definition any such optically active or racemic form which possesses the property of being an agonist or antagonist at the neuropeptide Y5
30 receptor. The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. Similarly, binding to the neuropeptide

Y5 receptor may be evaluated using the standard laboratory techniques referred to hereinafter.

The invention also relates to any and all tautomeric forms of the compounds of the formula (I) that possess neuropeptide Y5 receptor agonist or antagonist activity.

It will also be understood that certain compounds of the present invention may exist in solvated, for example hydrated, as well as unsolvated forms. It is to be understood that the present invention encompasses all such solvated forms which possess the property of interacting with the neuropeptide Y5 receptor.

In another aspect of the present invention there is provided a compound of formula (I'):



(I')

wherein:

R^1 is cyano, halo, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl, C_{1-4} alkoxy, $N-(C_{1-4}$ alkyl)amino or $N,N-(C_{1-4}$ alkyl)₂amino;

R^2 is halo, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl or C_{1-4} alkoxy;

A is $-NR^a-$, $-O-$ or a direct bond; wherein R^a is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl; wherein said C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl may be optionally substituted by one or more R^4 ;

R^3 is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl or C_{2-10} alkynyl wherein said C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl may be optionally substituted by one or more R^5 ; or R^3 is carbocyclyl or heterocyclyl wherein said carbocyclyl or heterocyclyl may be optionally substituted on carbon by one or more R^6 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R^7 ;

R^4 and R^5 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, C_{1-6} alkanoylamino, C_{2-6} alkenyloxy, carbonyl, C_{1-6} alkoxycarbonyl, heterocyclyl, heterocycliloxy, heterocyclylcarbonyl, heterocycliloxy, carbonyl, carbocyclyl, carbocycliloxy, carbocyclylcarbonyl, carbocycliloxy, carbonyl, $N-(C_{1-6}$ alkyl)amino, $N,N-(C_{1-6}$ alkyl)₂amino, C_{1-6} alkoxycarbonylamino, $N-(C_{1-6}$ alkyl)carbamoyl,

- 12 -

N,N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkylS(O)_a wherein a is 0-2, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl; wherein any C₁₋₆alkyl, heterocyclyl or carbocyclyl may be optionally substituted on carbon by one or more R⁶; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R⁷;

- 5 R⁶ is selected from halo, hydroxy, cyano, carbamoyl, ureido, trifluoromethyl, trifluoromethoxy, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy, C₁₋₄alkoxycarbonyl, C₂₋₄alkenyloxycarbonyl, C₁₋₄alkanoyl, C₁₋₄alkanoylamino, C₁₋₄alkanoyloxy, *N*-(C₁₋₄alkyl)amino, *N,N*-(C₁₋₄alkyl)₂amino, *N*-(C₁₋₄alkyl)carbamoyl, *N,N*-(C₁₋₄alkyl)₂carbamoyl, C₁₋₄alkylS(O)_a wherein a is 0-2,
- 10 *N*-(C₁₋₄alkyl)sulphamoyl, *N,N*-(C₁₋₄alkyl)₂sulphamoyl, heterocyclyl, heterocyclyloxy, heterocyclylcarbonyl, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylcarbonyl and carbocyclyloxycarbonyl; wherein any C₁₋₄alkyl, carbocyclyl and heterocyclyl may be optionally substituted on carbon by one or more R⁸;

- R⁷ is selected from C₁₋₄alkyl, C₁₋₄alkanoyl, C₁₋₄alkylsulphonyl, C₁₋₄alkoxycarbonyl,
- 15 carbamoyl, *N*-(C₁₋₄alkyl)carbamoyl, *N,N*-(C₁₋₄alkyl)₂carbamoyl, benzyl, benzoyl, phenylsulphonyl and phenyl;

- R⁸ is selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, methoxy, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, *N*-methylcarbamoyl,
- 20 *N,N*-dimethylcarbamoyl, methylthio, methylsulphanyl, mesyl, *N*-methylsulphamoyl, *N,N*-dimethylsulphamoyl, heterocyclyl, heterocyclyloxy, heterocyclylcarbonyl, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylcarbonyl and carbocyclyloxycarbonyl;

- x is 0 - 4; wherein the values of R¹ may be the same or different;
- 25 y is 0 - 3; wherein the values of R² may be the same or different; and
- n is 0 - 2;

or a pharmaceutically acceptable salt, prodrug or solvate thereof;
with the provisos:

- 1) when R³ is a nitrogen linked heterocyclyl, A is a direct bond;
- 30 2) when x and y are 0 and n is 2, the group R³-A-C(O)-NH- is not 2-formamido, 2-acetamido, 3-acetamido, 2-propionamido, 3-[2-(fur-2-ylcarbonylmethyl)acetamido], 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido),

- 2-(2-pthalimido-2-isopropylacetamido), 2-(2-pthalimido-2-isobutylacetamido),
 2-(2-pthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido),
 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido),
 2-(2-amino-2-benzylacetamido), 3-(2-pthalimidoacetamido), 3-(3-pthalimidopropionamido),
 5 3-(2-pthalimido-2-isopropylacetamido), 3-(2-pthalimido-2-isobutylacetamido),
 3-(2-pthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido),
 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-benzylacetamido) or
 2-(3,3,3-trifluoro-2-hydroxy-2-methylpropionamido);
 3) when x and y are 0 and n is 1, the group $R^3-A-C(O)-NH-$ is not 3-acetamido,
 10 3-(2-pthalimidoacetamido), 3-(3-pthalimidopropionamido),
 3-(2-pthalimido-2-isopropylacetamido), 3-(2-pthalimido-2-isobutylacetamido),
 3-(2-pthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido),
 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-isobutylacetamido) or
 3-(2-amino-2-benzylacetamido);
 15 4) when x and y are 0 and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-benzamido, 2-acetamido,
 2-benzyloxycarbonylamino, 2-(2-pthalimidoacetamido), 2-(3-pthalimidopropionamido),
 2-(2-pthalimido-2-isopropylacetamido), 2-(2-pthalimido-2-isobutylacetamido),
 2-(2-pthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido),
 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido),
 20 2-(2-amino-2-benzylacetamido), 3-(trifluoroacetamido), 3-benzyloxycarbonylamino or
 4-t-butyloxycarbonylamino;
 5) when $(R^1)_x$ is 7-fluoro, y is 0 and n is 2, the group $R^3-A-C(O)-NH-$ is not 3-acetamido;
 6) when x is 0, $(R^2)_y$ is 1-cyano and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido;
 7) when x is 0, $(R^2)_y$ is 3-bromo and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido or
 25 2-benzamido;
 8) when x is 0, $(R^2)_y$ is 1-bromo and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido or
 2-benzamido;
 9) when x is 0, $(R^2)_y$ is 1-chloro or 4-chloro and n is 0, the group $R^3-A-C(O)-NH-$ is not
 2-acetamido.
 30 Preferred values of R^1 , R^2 , R^3 , R^4 , X, A, x, y, z and n are as follows. Such values may
 be used where appropriate with any of the definitions, claims or embodiments defined
 hereinbefore or hereinafter. For the avoidance of doubt R^1 , R^2 , R^3 , R^4 , X, A, x, y, z and n as

used below correspond to those used in formula (I). It is to be understood that for compounds of formula (I'), (IA) and (IB) R^4 and z are irrelevant, R^5 as used in this section corresponds to R^4 ; R^6 as used in this section corresponds to R^5 ; R^7 as used in this section corresponds to R^6 ; R^8 as used in this section corresponds to R^7 ; and R^{11} as used in this section corresponds to R^8 .

5 In one aspect of the invention, preferably x is 0-3; wherein the values of R^1 may be the same or different.

In a further aspect of the invention, preferably x is 0-2; wherein the values of R^1 may be the same or different.

In an additional aspect of the invention, preferably x is 0-1.

10 Preferably x is 0.

In one aspect of the invention, preferably y is 0-2; wherein the values of R^2 may be the same or different.

In an additional aspect of the invention, preferably y is 0-1.

Preferably y is 0.

15 In one aspect of the invention, preferably X is a group of formula (A) (as depicted above).

In another aspect of the invention, preferably X is a group of formula (B) (as depicted above).

Preferably X is in the 2-position of the dibenzothiophene ring.

20 Preferably R^1 is halo or C_{1-4} alkyl.

More preferably R^1 is fluoro or methyl.

Preferably R^2 is halo, cyano or C_{1-4} alkyl.

More preferably R^2 is bromo, cyano or methyl.

Particularly R^2 is 3-bromo, 1-cyano, 1-methyl or 3-methyl.

25 In one aspect of the invention, preferably A is $—NR^a—$.

In another aspect of the invention, preferably A is $—O—$

In a further aspect of the invention, preferably A is a direct bond.

Preferably A is $—NR^a—$ or a direct bond.

30 one or more R^5 .
Preferably when A is $—NR^a—$; R^a is hydrogen or C_{1-10} alkyl optionally substituted by

More preferably when A is $—NR^a—$; R^a is hydrogen or unsubstituted C_{1-10} alkyl.

Particularly when A is $—NR^a—$; R^a is hydrogen or unsubstituted C_{1-4} alkyl.

- 15 -

More particularly when A is $\text{---NR}^a\text{---}$; R^a is methyl.

In one aspect of the invention, more particularly when A is $\text{---NR}^a\text{---}$, R^a is hydrogen.

In another aspect of the invention, more particularly when A is $\text{---NR}^a\text{---}$; R^a is C_{1-4} alkyl.

5 Preferably A is $\text{---NR}^a\text{---}$, ---O--- or a direct bond; wherein R^a is hydrogen, C_{1-10} alkyl or C_{2-10} alkenyl; wherein R^a may be optionally substituted by one or more R^5 ; wherein

R^3 is C_{1-10} alkyl or C_{2-10} alkenyl wherein R^3 may be optionally substituted by one or more R^6 ; or R^3 is carbocyclyl or heterocyclyl wherein R^3 may be optionally substituted on carbon by one or more R^7 ; and wherein if said heterocyclyl contains an -NH- moiety that
10 nitrogen may be optionally substituted by R^8 ;

R^5 and R^6 are independently selected from hydroxy, cyano, amino, C_{1-6} alkoxy, heterocyclyl, heterocycliloxy, carbocyclyl, carbocycliloxy, carbocyclyl-*N*-(C_{1-6} alkyl)amino, *N,N*-(C_{1-6} alkyl)₂amino, C_{1-6} alkoxycarbonylamino, C_{1-6} alkylS(O)_{*a*} wherein *a* is 0-2, *N,N*-(C_{1-6} alkyl)₂sulphamoylamino and C_{1-6} alkylsulphonylamino; wherein R^5 and R^6 may be
15 independently optionally substituted on carbon by one or more R^9 ;

R^7 and R^9 are independently selected from halo, hydroxy, nitro, carboxy, carbamoyl, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, C_{1-4} alkanoyl, C_{1-4} alkanoylamino, *N,N*-(C_{1-4} alkyl)₂amino, *N,N*-(C_{1-4} alkyl)₂carbamoyl, heterocyclyl, heterocyclylcarbonyl, carbocyclyl and carbocyclylcarbonylamino; wherein R^7 and R^9 may be independently
20 optionally substituted on carbon by one or more R^{11} ;

R^8 is selected from C_{1-4} alkyl, C_{1-4} alkylsulphonyl, *N,N*-(C_{1-4} alkyl)₂sulphamoyl, heterocyclyl and carbocyclyl; wherein R^8 may be optionally substituted on carbon by one or more R^{12} ;

R^{11} and R^{12} are independently selected from halo, hydroxy, cyano, carbamoyl, methyl, methoxy, allyloxy, heterocyclyl and carbocyclyl.
25

More preferably A is $\text{---NR}^a\text{---}$, ---O--- or a direct bond; wherein R^a is hydrogen, methyl, ethyl, propyl, isopropyl or allyl; wherein said methyl, ethyl, propyl, isopropyl or allyl may be optionally substituted by one or more R^5 ; wherein

R^3 is methyl, ethyl, propyl, butyl, hexyl or allyl wherein R^3 may be optionally
30 substituted by one or more R^6 ; or R^3 is selected from cyclopropyl, cyclopentyl, cyclohexyl, bicyclo[2,2,1]heptyl, phenyl, tetrahydropyranyl, morpholino, 2-oxopyrrolidinyl, piperidinyl, pyrrolidinyl, azetidiny, tetrahydrofuryl, 1,4-dioxo-8-azaspiro[4.5]decanyl,

- 16 -

1,1-dioxotetrahydrothienyl, 1,2,3,4,-tetrahydro-9H-pyrido[3,4-b]indole, decahydroquinolin-1-yl, 1,2,5,6-tetrahydropyridyl or piperazinyl; wherein R³ may be optionally substituted on carbon by one or more R⁷; and wherein if any heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R⁸;

5 R⁵ is selected from cyano, methoxy, pyridyl;

R⁶ is selected from hydroxy, cyano, amino, methoxy, pyridyl, 2-oxopyrrolidinyl, 1,2,4-triazolyl, 1,1-dioxotetrahydrothienyl, thienyl, 2-oxooxazolidinyl, imidazolyl, 1,1-dioxothiomorpholino, 2-oxo-1,2-dihydropyridyl, benzimidazolyl, pyrazolyl, succinimido, tetrahydrofuryl, 2,4-dioxothiazolidinyl, morpholino, furyl, pyridyloxy, pyridazinyl, 10 cyclopentyl, cyclohexyl, phenyl, phenoxy, *N*-methylanilino, *N,N*-dimethylamino, *t*-butoxycarbonylamino, mesyl, *N,N*-dimethylsulphamoylamino and isopropylsulphonylamino; wherein R⁶ may be optionally substituted on carbon by one or more R⁹;

R⁷ is selected from fluoro, hydroxy, nitro, carboxy, carbamoyl, methyl, ethyl, propyl, 15 methoxycarbonyl, ethoxycarbonyl, acetyl, acetamido, *N,N*-dimethylamino, *N,N*-dimethylcarbamoyl, *N,N*-diethylcarbamoyl, 1,4-dihydrooxazin-2-one, pyrazolyl, 2-oxo-2,3-dihydrobenzimidazolyl, pyridyl, 2,3-dihydro-2-oxobenzimidazolyl, morpholinocarbonyl, phenyl and cyclohexylcarbonylamino; wherein any R⁷ may be optionally substituted on carbon by one or more R¹¹;

20 R⁹ is selected from fluoro, chloro, hydroxy, methyl, methoxy, pyridyl and phenyl;

R⁸ is selected from methyl, ethyl, propyl, pentyl, ethylsulphonyl, *N,N*-dimethylsulphamoyl, thieno[2,3-d]pyrimidinyl, pyrimidinyl, pyridyl and phenyl; wherein any R⁸ may be optionally substituted on carbon by one or more R¹²;

R¹¹ is selected from hydroxy, cyano, allyloxy, pyrrolidinyl, furyl and phenyl;

25 R¹² is selected from fluoro, chloro, hydroxy, carbamoyl, methyl, methoxy, tetrahydrofuryl, morpholino and phenyl.

Preferably R³ is C₁₋₁₀alkyl optionally substituted by one or more R⁶; or R³ is carbocyclyl or heterocyclyl wherein said carbocyclyl or heterocyclyl may be optionally substituted on carbon by one or more R⁷; and wherein if said heterocyclyl contains an -NH- 30 moiety that nitrogen may be optionally substituted by R⁸.

More preferably R³ is C₁₋₄alkyl optionally substituted by one or more R⁶; or R³ is carbocyclyl or heterocyclyl wherein said carbocyclyl or heterocyclyl may be optionally

- 17 -

substituted on carbon by one or more R⁷; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R⁸.

Particularly R³ is methyl, ethyl, isopropyl or t-butyl; wherein said methyl, ethyl, isopropyl or t-butyl may be optionally substituted by one or more R⁶; or R³ is cyclopropyl, phenyl, morpholino, piperazin-1-yl, tetrahydropyran-4-yl, piperid-1-yl, piperid-4-yl; wherein said cyclopropyl, phenyl, morpholino, piperazin-1-yl, tetrahydropyran-4-yl, piperid-1-yl or piperid-4-yl may be optionally substituted on carbon by one or more R⁷; and wherein piperazin-1-yl and piperid-4-yl may be optionally substituted on nitrogen by R⁸.

More particularly R³ is methyl, ethyl, isopropyl or t-butyl; wherein said methyl, ethyl, isopropyl or t-butyl may be optionally substituted by one or more fluoro, hydroxy, cyano, amino, methoxy, pyridyl, morpholino, 2-pyrrolidinonyl, triazolyl, 1,1-dioxotetrahydrothienyl, 2-oxazolidinonyl, phenoxy, pyridyloxy, imidazolyl, 1,1-dioxothiomorpholino, *N,N*-dimethylamino, t-butyloxycarbonylamino; or R³ is cyclopropyl, 4-nitrophenyl, morpholino, 4-methylpiperazin-1-yl, tetrahydropyran-4-yl, 4-hydroxymethylpiperid-1-yl or 1-methylpiperid-4-yl.

Particularly preferred R³ is methyl, cyanomethyl, 2-pyrrolidinon-1-ylmethyl, 1,1-dioxothiomorpholinomethyl, pyrid-3-yloxymethyl, 1,1-dioxotetrahydrothien-3-ylmethyl, 2-oxazolidinon-3-ylmethyl, pyrid-4-ylmethyl, 1,2,4-triazol-1-ylmethyl, 1-phenoxyethyl, 2-imidazol-1-ylethyl, 2-*N,N*-dimethylaminoethyl, 2-1,2,4-triazol-1-ylethyl, 2-methoxyethyl, 2-pyrid-4-ylethyl, isopropyl, 1-pyrid-4-ylprop-2-yl, 2-aminoprop-2-yl, 2-hydroxy-3,3,3-trifluoroprop-2-yl, 1-morpholinoprop-2-yl, 2-(t-butoxycarbonylamino)prop-2-yl, t-butyl, cyclopropyl, 4-nitrophenyl, morpholino, 4-methylpiperazin-1-yl, tetrahydropyran-4-yl, 4-hydroxymethylpiperid-1-yl or 1-methylpiperid-4-yl.

More particularly preferred R³ is 1,1-dioxotetrahydrothien-3-ylmethyl, 1-phenoxyethyl, 2-pyrid-4-ylethyl, isopropyl, 1-pyrid-4-ylprop-2-yl, 1-morpholinoprop-2-yl, t-butyl, tetrahydropyran-4-yl or 1-methylpiperid-4-yl.

Preferably the group R³-A- is methyl, 2-oxo-pyrrolidin-1-ylmethyl, 1,2,4-triazol-1-ylmethyl, 1,1-dioxotetrahydrothien-3-ylmethyl, 2-oxooxazolidin-3-ylmethyl, pyrid-3-yloxymethyl, 1,1-dioxothiomorpholinomethyl, cyanomethyl, 2-oxo-1,2-dihydropyrid-1-ylmethyl, 2-oxocyclopentylmethyl, succinimidomethyl, 3-benzyl-2-oxopyrrolidin-1-ylmethyl, 3-hydroxypyridazin-6-yloxymethyl, 2-pyrid-4-ylethyl,

- 2-methoxyethyl, 1-phenoxyethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-imidazol-1-ylethyl,
succinimidoethyl, prop-2-yl, 3,3,3-trifluoro-2-hydroxyprop-2-yl, 1-morpholinoprop-2-yl,
1-pyrid-4-ylprop-2-yl, 2-aminoprop-2-yl, 2-(*t*-butoxycarbonylamino)prop-2-yl,
1-(1,2,4-triazol-1-yl)prop-2-yl, 2-pyrid-4-ylpropyl, *t*-butyl, 1,1,1-trifluorobut-3-yl,
5 1-hydroxyhex-2-yl, cyclopropyl, 3-hydroxybicyclo[2.2.1]hept-2-yl, 4-nitrophenyl,
morpholino, 4-methylpiperazin-1-yl, tetrahydropyran-4-yl, 4-hydroxymethylpiperidin-1-yl,
1-methyl-2-oxopyrrolidin-4-yl, 2-(pyrrolidin-1-ylmethyl)pyrrolidinyl,
3-carbamoylpiperidin-1-yl, 3-hydroxyazetidin-1-yl, 2-(allyloxymethyl)morpholino,
4-(1,4-dihydrooxazin-2-one-3-yl)piperidin-1-yl, 4-(*N,N*-dimethylsulphamoyl)piperazin-1-yl,
10 4-hydroxyethylpiperidin-1-yl, 4-(tetrahydrofur-2-ylmethyl)piperazin-1-yl,
4-(3-methoxypropyl)piperazin-1-yl, 4-pyrid-4-ylpiperidin-1-yl, 4-pyrid-2-ylpiperazin-1-yl,
3-(*N,N*-dimethylamino)pyrrolidin-1-yl, 4-carboxypiperidin-1-yl,
1-methyl-2-oxo-5-phenylpyrrolidin-4-yl, 2-oxo-5,5-dimethyltetrahydrofur-4-yl,
tetrahydrofur-4-yl, 2,2-dimethyltetrahydropyran-4-yl, 1-benzyl-2-oxopyrrolidin-4-yl,
15 2-oxo-5-phenyltetrahydrofuryl, 2-(3-hydroxypropyl)piperidin-1-yl,
4-(2-carbamoylethyl)piperazin-1-yl, 3-oxo-4-(2-methoxyethyl)piperazin-1-yl,
4-(*N,N*-dimethylamino)-4-carbamoylpiperidin-1-yl, 4-(2-morpholinoethyl)piperazin-1-yl,
1,4-dioxo-8-azaspiro[4.5]decan-8-yl, 1,1-dioxotetrahydrothien-3-yl,
4-ethylsulphonylpiperazin-1-yl, 4-(thieno[2,3-*d*]pyrimidin-4-yl)piperazin-1-yl,
20 4-(2-hydroxyethyl)piperazin-1-yl, 4-carbamoylpiperidin-1-yl,
3-methyl-3-phenylpiperidin-1-yl, 2-benzyloxycarbonylpiperidin-1-yl,
4-(*N,N*-dimethylcarbamoyl)piperidin-1-yl, 3-(pyrid-4-yl)pyrrolidin-1-yl,
3-(pyrid-3-yl)pyrrolidin-1-yl, 4,4-difluoropiperidin-1-yl,
4-(2-methylpyrimidin-4-yl)piperazin-1-yl, 4-(2,3,5,6-tetrafluoropyrid-4-yl)piperazin-1-yl,
25 4-(pyrimidin-4-yl)piperazin-1-yl, 3-cyanomethylpiperidin-1-yl,
4-cyclohexylcarbonylaminopiperidin-1-yl, 4-acetamidopiperidin-1-yl,
4-(6-chloropyrimidin-4-yl)piperazin-1-yl, 4-(pent-3-yl)piperazin-1-yl,
1,2,5,6-tetrahydropyrid-1-yl, 1-methylpiperidin-1-yl, 2-methylpiperidin-1-yl,
decahydroquinolin-1-yl, 3-ethoxycarbonyl-4-oxopiperidin-1-yl,
30 2-ethoxycarbonylpyrrolidin-1-yl, 4-acetylpiperidin-1-yl, 2-azabicyclo[2.2.1]hept-2-yl,
1,2,3,4,-tetrahydro-9H-pyrido[3,4-*b*]indol-2-yl,
4-(2,3-dihydro-2-oxobenzimidazol-1-yl)piperidin-1-yl, 4-(2-methoxyphenyl)piperazin-1-yl,

- 4-methylpiperidin-1-yl, 4-(3-chlorophenyl)piperazin-1-yl,
4-(3-methoxyphenyl)piperazin-1-yl, 4-(4-methoxyphenyl)piperazin-1-yl,
4-(4-chlorophenyl)piperazin-1-yl,
4-(2,3-dihydro-2-oxobenzimidazol-1-yl)-1,2,5,6-tetrahydropyrid-1-yl, 4-phenylpiperidin-1-yl,
5 4-(3-fur-2-ylpyrazol-5-yl)piperidin-1-yl, 2-pyrid-4-ylethylamino,
3-imidazol-1-ylpropylamino, 4-hydroxycyclohexylamino,
2-(*N,N*-dimethylsulphamoylamino)ethylamino, 2-(isopropylsulphonylamino)ethylamino,
2-imidazol-5-ylethylamino, 2-mesyethylamino, 2-morpholinoethylamino,
1-methoxycarbonylcyclopropylamino, 1-benzylpyrrolidin-3-ylamino,
10 3-(*N*-methylanilino)propylamino, 2-(5-methyl-2,4-dioxothiazolidin-3-yl)ethylamino,
2-(*t*-butoxycarbonylamino)ethylamino, *N*-(*N*-methyl-*N*-pyrid-3-ylmethylaminopropyl)amino,
1-cyclohexylethylamino, *N*-methyl-*N*-(2-pyrid-4-ylethyl)amino,
N-methyl-*N*-(2-pyrid-2-ylethyl)amino, *N*-methyl-*N*-(2-cyanoethyl)amino,
N-methyl-*N*-(pyrid-3-ylmethyl)amino, *N*-methyl-*N*-(2-*N,N*-dimethylaminoethyl)amino,
15 *N*-methyl-*N*-(1-methylpiperidin-4-yl)amino, *N*-methyl-*N*-(3-mesylpropyl)amino,
N-methyl-*N*-(4-hydroxy-4-methyltetrahydropyran-3-yl)amino,
N-(pyrid-3-ylmethyl)-*N*-(2-cyanoethyl)amino, *N*-methyl-*N*-(2-hydroxypropyl)amino,
N-methyl-*N*-(2,2-dimethoxyethyl)amino, *N*-methyl-*N*-phenethylamino,
N-methyl-*N*-(tetrahydrofur-2-ylmethyl)amino, *N*-methyl-*N*-(2-morpholinoethyl)amino,
20 *N*-methyl-*N*-(6-methylpyrid-2-ylmethyl)amino, *N*-methyl-*N*-(1-methylpyrrolidin-3-yl)amino,
N-methyl-*N*-[2-(4-hydroxyphenyl)-2-hydroxyethyl]amino,
N-methyl-*N*-(1-benzylpyrrolidin-3-yl)amino, *N*-methyl-*N*-[2-(1,2,4-triazol-1-yl)ethyl]amino,
N-methyl-*N*-(fur-2-ylmethyl)amino, *N*-methyl-*N*-(benzimidazol-2-ylmethyl)amino,
N-methyl-*N*-benzylamino, *N*-methyl-*N*-(2-chlorobenzyl)amino,
25 *N*-methyl-*N*-(3-chlorobenzyl)amino, *N*-methyl-*N*-(4-chlorobenzyl)amino,
N-methyl-*N*-[2-(3,4-dimethoxypyrid-4-yl)ethyl]amino,
N-methyl-*N*-(5-phenylpyrazol-3-ylmethyl)amino, *N*-methyl-*N*-(4-fluorobenzyl)amino,
N-methyl-*N*-(2-methoxyphenylprop-2-yl)amino, *N*-ethyl-*N*-(pyrid-4-ylmethyl)amino,
N-(2-methoxyethyl)-*N*-(pyrid-3-ylmethyl)amino, *N*-ethyl-*N*-(2-methoxyethyl)amino,
30 *N*-(2-hydroxyethyl)-*N*-isopropylamino, *N*-(2-cyanoethyl)-*N*-(3-morpholinopropyl)amino,
N-(2-cyanoethyl)-*N*-(thien-2-ylmethyl)amino, *N*-(2-cyanoethyl)-*N*-benzylamino,
N-ethyl-*N*-(1-benzylpyrrolidin-3-yl)amino, *N*-ethyl-*N*-(1,1-dioxotetrahydrothien-3-yl)amino,

N-(2-pyrid-4-ylethyl)-*N*-cyclopropylamino,
N-(2-hydroxy-2-pyrid-4-ylethyl)-*N*-isopropylamino,
N-(4-hydroxycyclohexyl)-*N*-isopropylamino, *N*-allyl-*N*-(1,1-dioxotetrahydrothien-3-yl)amino,
diallylamino, *N*-allyl-*N*-cyclopentylamino or *N*-benzyl-*N*-propylamino.

- 5 More preferably the group R³-A- is 1,1,1-trifluorobut-3-yl, 4-pyrid-4-ylpiperidin-1-yl,
4-pyrid-2-ylpiperidin-1-yl, 4-(2,3,5,6-tetrafluoropyrid-4-yl)piperazin-1-yl,
N-methyl-*N*-(2-pyrid-4-ylethyl)amino or *N*-methyl-*N*-phenethylamino.

Preferably R⁵ and R⁶ are independently selected from halo, hydroxy, cyano, amino,
C₁₋₆alkoxy, heterocyclyl, heterocycloxy, carbocycloxy, *N,N*-(C₁₋₆alkyl)₂amino,

- 10 C₁₋₆alkoxycarbonylamino.

More preferably R⁵ and R⁶ are independently selected from fluoro, hydroxy, cyano,
amino, methoxy, pyridyl, morpholino, 2-pyrrolidinonyl, triazolyl, 1,1-dioxotetrahydrothienyl,
2-oxazolidinonyl, phenoxy, pyridyloxy, imidazolyl, 1,1-dioxothiomorpholino,
N,N-dimethylamino, *t*-butyloxycarbonylamino.

- 15 Particularly R⁵ and R⁶ are independently selected from pyridyl, morpholino,
1,1-dioxotetrahydrothienyl or phenoxy.

Preferably R⁷ is hydroxymethyl.

Preferably R⁸ is selected from methyl.

Preferably R¹¹ is hydroxy.

- 20 In one aspect of the invention, preferably *n* is 0.

In another aspect of the invention, preferably *n* is 1.

In a further aspect of the invention preferably *n* is 2.

Preferably R⁴ is C₁₋₄alkyl.

More preferably R⁴ is methyl.

- 25 Particularly R⁴ is 6-methyl.

In one aspect of the invention, preferably *z* is 0-2; wherein the values of R⁴ may be the
same or different.

In an additional aspect of the invention, preferably *z* is 0-1.

In another aspect of the invention, preferably *z* is 1.

- 30 In a further additional aspect of the invention, preferably *z* is 0.

In one aspect of the invention, preferably *n* is 2.

In an additional aspect of the invention, preferably *n* is 1.

- 21 -

In a further additional aspect of the invention, preferably n is 0.

Therefore in one aspect of the invention, there is provided a compound of formula (I) as depicted above wherein:

x is 0;

5 y is 0;

A is $\text{—NR}^a\text{—}$, —O— or a direct bond; wherein R^a is hydrogen or C_{1-4} alkyl;

R^3 is C_{1-4} alkyl optionally substituted by one or more R^6 ; or R^3 is carbocyclyl or heterocyclyl wherein said carbocyclyl or heterocyclyl may be optionally substituted on carbon by one or more R^7 ; and wherein if said heterocyclyl contains an —NH— moiety that nitrogen
10 may be optionally substituted by R^8 ;

R^6 is selected from halo, hydroxy, cyano, amino, C_{1-6} alkoxy, heterocyclyl, heterocyclylloxy, carbocyclylloxy, $\text{N,N-(C}_{1-6}\text{alkyl)}_2$ amino, C_{1-6} alkoxycarbonylamino;

R^7 is hydroxymethyl;

R^8 is methyl; and

15 n is 0 - 2;

or a pharmaceutically acceptable salt, prodrug or solvate thereof;

with the provisos:

- 1) when R^3 is a nitrogen linked heterocyclyl, A is a direct bond;
- 2) when n is 2, the group $\text{R}^3\text{—A—C(O)—NH—}$ is not 2-acetamido, 3-acetamido, 2-propionamido,
20 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido),
2-(2-phthalimido-2-isopropylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido),
2-(2-amino-2-isopropylacetamido), 3-(2-phthalimidoacetamido),
3-(3-phthalimidopropionamido), 3-(2-phthalimido-2-isopropylacetamido),
3-(2-aminoacetamido), 3-(3-aminopropionamido), 3-(2-amino-2-isopropylacetamido) or
25 2-(3,3,3-trifluoro-2-hydroxy-2-methylpropionamido);
- 3) when n is 1, the group $\text{R}^3\text{—A—C(O)—NH—}$ is not 3-acetamido, 3-(2-phthalimidoacetamido),
3-(3-phthalimidopropionamido), 3-(2-phthalimido-2-isopropylacetamido),
3-(2-aminoacetamido), 3-(3-aminopropionamido) or 3-(2-amino-2-isopropylacetamido);
- 4) when n is 0, the group $\text{R}^3\text{—A—C(O)—NH—}$ is not 2-acetamido, 2-benzyloxycarbonylamino,
30 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido),
2-(2-phthalimido-2-isopropylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido),

- 22 -

2-(2-amino-2-isopropylacetamido), 3-(trifluoroacetamido), 3-benzyloxycarbonylamino or 4-t-butyloxycarbonylamino.

Therefore in a further aspect of the invention, there is provided a compound of formula (I) as depicted above wherein:

5 x is 0;

y is 0;

A is $\text{---NR}^a\text{---}$, ---O--- or a direct bond; wherein R^a is hydrogen or methyl;

R^3 is methyl, cyanomethyl, 2-pyrrolidinon-1-ylmethyl,

1,1-dioxothiomorpholinomethyl, pyrid-3-ylloxymethyl, 1,1-dioxotetrahydrothien-3-ylmethyl,

10 2-oxazolidinon-3-ylmethyl, pyrid-4-ylmethyl, 1,2,4-triazol-1-ylmethyl, 1-phenoxyethyl,

2-imidazol-1-ylethyl, 2-*N,N*-dimethylaminoethyl, 2-1,2,4-triazol-1-ylethyl, 2-methoxyethyl,

2-pyrid-4-ylethyl, isopropyl, 1-pyrid-4-ylprop-2-yl, 2-aminoprop-2-yl,

2-hydroxy-3,3,3-trifluoroprop-2-yl, 1-morpholinoprop-2-yl,

2-(t-butoxycarbonylamino)prop-2-yl, t-butyl, cyclopropyl, 4-nitrophenyl, morpholino,

15 4-methylpiperazin-1-yl, tetrahydropyran-4-yl, 4-hydroxymethylpiperid-1-yl or

1-methylpiperid-4-yl; and

n is 0 - 2;

or a pharmaceutically acceptable salt, prodrug or solvate thereof;

with the provisos:

20 1) when R^3 is morpholino, 4-methylpiperazin-1-yl or 4-hydroxymethylpiperid-1-yl, A is a direct bond;

2) when n is 2, the group $R^3\text{-A-C(O)-NH-}$ is not 2-acetamido, 3-acetamido, or 2-(3,3,3-trifluoro-2-hydroxy-2-methylpropionamido);

3) when n is 1, the group $R^3\text{-A-C(O)-NH-}$ is not 3-acetamido;

25 4) when n is 0, the group $R^3\text{-A-C(O)-NH-}$ is not 2-acetamido.

Therefore in an additional aspect of the invention, there is provided a compound of formula (I) as depicted above wherein:

x is 0;

y is 0;

30 A is $\text{---NR}^a\text{---}$ or a direct bond; wherein R^a is methyl;

R³ is 1,1-dioxotetrahydrothien-3-ylmethyl, 1-phenoxyethyl, 2-pyrid-4-ylethyl, isopropyl, 1-pyrid-4-ylprop-2-yl, 1-morpholinoprop-2-yl, t-butyl, tetrahydropyran-4-yl or 1-methylpiperid-4-yl; and

n is 0 - 2;

5 or a pharmaceutically acceptable salt, prodrug or solvate thereof.

Therefore in another aspect of the invention there is provided a compound of formula (I) (as depicted above) wherein:

X is a group of formula (A) (as depicted above) or a group of formula (B) (as depicted above);

10 R² is halo, cyano or C₁₋₄alkyl;

A is —NR^a—, —O— or a direct bond; wherein R^a is hydrogen, C₁₋₁₀alkyl or C₂₋₁₀alkenyl; wherein R^a may be optionally substituted by one or more R⁵;

R³ is C₁₋₁₀alkyl or C₂₋₁₀alkenyl wherein R³ may be optionally substituted by one or more R⁶; or R³ is carbocyclyl or heterocyclyl wherein R³ may be optionally substituted on
15 carbon by one or more R⁷; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R⁸;

R⁵ and R⁶ are independently selected from hydroxy, cyano, amino, C₁₋₆alkoxy, heterocyclyl, heterocyclxyloxy, carbocyclyl, carbocyclxyloxy, carbocyclyl-*N*-(C₁₋₆alkyl)amino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkoxycarbonylamino, C₁₋₆alkylS(O)_a wherein a is 0-2,
20 *N,N*-(C₁₋₆alkyl)₂sulphamoylamino and C₁₋₆alkylsulphonylamino; wherein R⁵ and R⁶ may be independently optionally substituted on carbon by one or more R⁹;

R⁷ and R⁹ are independently selected from halo, hydroxy, nitro, carboxy, carbamoyl, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxycarbonyl, C₁₋₄alkanoyl, C₁₋₄alkanoylamino, *N,N*-(C₁₋₄alkyl)₂amino, *N,N*-(C₁₋₄alkyl)₂carbamoyl, heterocyclyl, heterocyclylcarbonyl,
25 carbocyclyl and carbocyclylcarbonylamino; wherein R⁷ and R⁹ may be independently optionally substituted on carbon by one or more R¹¹;

R⁸ is selected from C₁₋₄alkyl, C₁₋₄alkylsulphonyl, *N,N*-(C₁₋₄alkyl)₂sulphamoyl, heterocyclyl and carbocyclyl; wherein R⁸ may be optionally substituted on carbon by one or more R¹²;

30 R¹¹ and R¹² are independently selected from halo, hydroxy, cyano, carbamoyl, methyl, methoxy, allyloxy, heterocyclyl and carbocyclyl;

R⁴ is C₁₋₄alkyl;

- 24 -

x is 0;

y is 0-1;

z is 1; and

n is 0-2;

- 5 or a pharmaceutically acceptable salt, prodrug or solvate thereof;
with the provisos when X is a group of formula (A):
- 1) when R^3 is a nitrogen linked heterocyclyl, A is a direct bond;
 - 2) when y is 0 and n is 2, the group $R^3-A-C(O)-NH-$ is not 2-acetamido, 3-acetamido, 2-propionamido, 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido),
 - 10 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido), 2-(2-phthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido), 2-(2-amino-2-benzylacetamido), 3-(2-phthalimidoacetamido), 3-(3-phthalimidopropionamido), 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido),
 - 15 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido), 3-(2-amino-2-isopropylacetamido) or 3-(2-amino-2-benzylacetamido);
 - 3) when y is 0 and n is 1, the group $R^3-A-C(O)-NH-$ is not 3-acetamido, 3-(2-phthalimidoacetamido), 3-(3-phthalimidopropionamido), 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido),
 - 20 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido), 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-isobutylacetamido) or 3-(2-amino-2-benzylacetamido);
 - 4) when y is 0 and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-benzamido, 2-acetamido, 2-benzyloxycarbonylamino, 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido),
 - 25 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido), 2-(2-phthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido), 2-(2-amino-2-benzylacetamido), 3-benzyloxycarbonylamino or 4-t-butyloxycarbonylamino;
 - 5) when $(R^2)_y$ is 1-cyano and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido;
 - 30 6) when $(R^2)_y$ is 3-bromo and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido or 2-benzamido;

7) when (R²)_y is 1-bromo and n is 0, the group R³-A-C(O)-NH- is not 2-acetamido or 2-benzamido; and

8) when (R²)_y is 1-chloro or 4-chloro and n is 0, the group R³-A-C(O)-NH- is not 2-acetamido.

Therefore in a further aspect of the invention there is provided a compound of formula

5 (I) (as depicted above) wherein:

X is a group of formula (A) (as depicted above) or a group of formula (B) (as depicted above) in the 2-position of the dibenzothiophene ring;

R² is bromo, cyano or methyl;

A is —NR^a—, —O— or a direct bond; wherein R^a is hydrogen, methyl, ethyl, propyl,
10 isopropyl or allyl; wherein said methyl, ethyl, propyl, isopropyl or allyl may be optionally substituted by one or more R⁵;

R³ is methyl, ethyl, propyl, butyl, hexyl or allyl wherein R³ may be optionally substituted by one or more R⁶; or R³ is selected from cyclopropyl, cyclopentyl, cyclohexyl, bicyclo[2,2,1]heptyl, phenyl, tetrahydropyranyl, morpholino, 2-oxopyrrolidinyl, piperidinyl,
15 pyrrolidinyl, azetidiny, tetrahydrofuryl, 1,4-dioxo-8-azaspiro[4.5]decanyl, 1,1-dioxotetrahydrothienyl, 1,2,3,4,-tetrahydro-9H-pyrido[3,4-b]indole, decahydroquinolin-1-yl, 1,2,5,6-tetrahydropyridyl or piperazinyl; wherein R³ may be optionally substituted on carbon by one or more R⁷; and wherein if any heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R⁸;

20 R⁵ is selected from cyano, methoxy, pyridyl;

R⁶ is selected from hydroxy, cyano, amino, methoxy, pyridyl, 2-oxopyrrolidinyl, 1,2,4-triazolyl, 1,1-dioxotetrahydrothienyl, thienyl, 2-oxooxazolidinyl, imidazolyl, 1,1-dioxothiomorpholino, 2-oxo-1,2-dihydropyridyl, benzimidazolyl, pyrazolyl, succinimido, tetrahydrofuryl, 2,4-dioxothiazolidinyl, morpholino, furyl, pyridyloxy, pyridazinyloxy,
25 cyclopentyl, cyclohexyl, phenyl, phenoxy, *N*-methylanilino, *N,N*-dimethylamino, *t*-butoxycarbonylamino, mesyl, *N,N*-dimethylsulphamoylamino and isopropylsulphonylamino; wherein R⁶ may be optionally substituted on carbon by one or more R⁹;

R⁷ is selected from fluoro, hydroxy, nitro, carboxy, carbamoyl, methyl, ethyl, propyl,
30 methoxycarbonyl, ethoxycarbonyl, acetyl, acetamido, *N,N*-dimethylamino, *N,N*-dimethylcarbamoyl, *N,N*-diethylcarbamoyl, 1,4-dihydrooxazin-2-one, pyrazolyl, 2-oxo-2,3-dihydrobenzimidazolyl, pyridyl, 2,3-dihydro-2-oxobenzimidazolyl,

- 26 -

morpholinocarbonyl, phenyl and cyclohexylcarbonylamino; wherein any R⁷ may be optionally substituted on carbon by one or more R¹¹;

R⁹ is selected from fluoro, chloro, hydroxy, methyl, methoxy, pyridyl and phenyl;

R⁸ is selected from methyl, ethyl, propyl, pentyl, ethylsulphonyl,

5 *N,N*-dimethylsulphamoyl, thieno[2,3-d]pyrimidinyl, pyrimidinyl, pyridyl and phenyl; wherein any R⁸ may be optionally substituted on carbon by one or more R¹²;

R¹¹ is selected from hydroxy, cyano, allyloxy, pyrrolidinyl, furyl and phenyl;

R¹² is selected from fluoro, chloro, hydroxy, carbamoyl, methyl, methoxy, tetrahydrofuryl, morpholino and phenyl;

10 R⁴ is methyl;

R⁴ is C₁₋₄alkyl;

x is 0;

y is 0-1;

z is 1; and

15 n is 0-2;

or a pharmaceutically acceptable salt, prodrug or solvate thereof;

with the provisos when X is a group of formula (A):

1) when R³ is a nitrogen linked heterocyclyl, A is a direct bond;

2) when y is 0 and n is 2, the group R³-A-C(O)-NH- is not 2-acetamido, 2-propionamido,
20 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido) or
2-(2-amino-2-benzylacetamido);

3) when y is 0 and n is 0, the group R³-A-C(O)-NH- is not 2-benzamido, 2-acetamido,
2-benzoyloxycarbonylamino, 2-(2-aminoacetamido), 2-(3-aminopropionamido),
2-(2-amino-2-isopropylacetamido) or 2-(2-amino-2-benzylacetamido);

25 4) when (R²)_y is 1-cyano and n is 0, the group R³-A-C(O)-NH- is not 2-acetamido;

5) when (R²)_y is 3-bromo and n is 0, the group R³-A-C(O)-NH- is not 2-acetamido or
2-benzamido; and

6) when (R²)_y is 1-bromo and n is 0, the group R³-A-C(O)-NH- is not 2-acetamido or
2-benzamido.

30 In another aspect of the invention, preferred compounds of the invention are any one
of the Examples or a pharmaceutically acceptable salt, prodrug or solvate thereof.

- 27 -


More preferred compounds of the invention are Examples 7, 19, 26, 27, 29, 30, 34, 35, 36, 37, 44, 45 and 47 or a pharmaceutically acceptable salt, prodrug or solvate thereof.

Further preferred compounds of the invention are Examples 19, 44, 50, 83, 105, 107 or 124 or a pharmaceutically acceptable salt, prodrug or solvate thereof.


5 Preferred aspects of the invention are those which relate to the compound of formula (I) or a pharmaceutically acceptable salt thereof.

Another aspect of the present invention provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof which process (wherein R^1 , R^2 , R^3 , A and n are, unless otherwise specified, as defined in
10 formula (I)) comprises of:

Process a): for compounds of formula (I) wherein A is a direct bond and X is a group of formula (A); reacting an amine of formula (II):




15 with an acid of formula (III):



or an activated derivative thereof; or

Process b): for compounds of formula (I) wherein $n > 0$; by oxidising a compound of formula
20 (I) where $n = 0$;

Process c): for compounds of formula (I) wherein A is $-NR^a-$ and X is a group of formula (A); by reacting a compound of formula (IV):



- 28 -


(IV)

wherein L is a displaceable group; with an amine of formula (V):




(V)

- 5 *Process d)*: for compounds of formula (I) wherein A is -NR^a- or -O- and X is a group of formula (A); reacting a compound of formula (II) with a compound of formula (VI):




(VI)

- Process e)*: for compounds of formula (I) wherein A is -NH- and X is a group of formula (A);
10 reacting a compound of formula (II) with an isocyanate of formula (VII):



(VII)

Process f): for compounds of formula (I) wherein X is a group of formula (B) reacting a compound of formula (II) with a compound of formula (VIII):



15

(VIII)

wherein L is a displaceable group;

and thereafter if necessary:

- i) converting a compound of the formula (I) into another compound of the formula (I);
20 ii) removing any protecting groups;
iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

L is a displaceable group. Suitable values for L are phenols for example p-nitrophenol or penta-fluorophenol.

Specific reaction conditions for the above reactions are as follows.

- 25 *Process a)* Amines of formula (II) and acids of formula (III) may be coupled together in the presence of a suitable coupling reagent. Standard peptide coupling reagents known in the art can be employed as suitable coupling reagents, or for example carbonyldiimidazole and dicyclohexyl-carbodiimide, optionally in the presence of a catalyst such as

dimethylaminopyridine or 4-pyrrolidinopyridine, optionally in the presence of a base for example triethylamine, pyridine, or 2,6-di-*alkyl*-pyridines such as 2,6-lutidine or 2,6-di-*tert*-butylpyridine. Suitable solvents include dimethylacetamide, dichloromethane, benzene, tetrahydrofuran and dimethylformamide. The coupling reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

Suitable activated acid derivatives include acid halides, for example acid chlorides, and active esters, for example pentafluorophenyl esters. The reaction of these types of compounds with amines is well known in the art, for example they may be reacted in the presence of a base, such as those described above, and in a suitable solvent, such as those described above. The reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

The amines of formula (II) and acids of formula (III) are commercially available or they are known compounds or they are prepared by processes known in the art.

Process b) Suitable oxidising agents include potassium permanganate, OXONE, sodium periodate, *tert*-butyl hydroperoxide (as solution in toluene), peracids (such as for example 3-chloroperoxybenzoic acid), hydrogen peroxide, TPAP (tetrapropylammonium perruthenate) or oxygen. The reaction may be conducted in a suitable solvent such as ether, dichloromethane, methanol, ethanol, water, acetic acid, or mixtures of two or more of these solvents. The reaction may conveniently be performed at a temperature in the range of -40 to 100°C.

Compounds of formula (I) where $n = 0$ may be prepared by processes a) or c).

Process c) Compounds of formula (IV) and amines of formula (V) may be reacted together in the presence of a suitable base, for example triethylamine, pyridine, or 2,6-di-*alkyl*-pyridines such as 2,6-lutidine or 2,6-di-*tert*-butylpyridine, or excess (V), in a suitable solvent such as dichloromethane, EtOAc or tetrahydrofuran. The reaction may conveniently be performed at a temperature in the range of -40 to 50°C.

The compounds of formula (IV) may be prepared from amines of formula (II) by standard processes known in the art. Compounds of formula (V) are commercially available or they are known compounds or they are prepared by processes known in the art.

Process d) Compounds of formula (II) and compounds of formula (VI) may be reacted in the presence of a base, such as those described above, in a suitable solvent, such as dichloromethane, toluene or tetrahydrofuran. The reaction may conveniently be performed at

a temperature in the range of -40 to 100°C.

Compounds of formula (VI) are commercially available or they are known compounds or they are prepared by processes known in the art.

Process e) Compounds of formula (II) and compounds of formula (VII) may be reacted in the presence of a suitable solvent, such as toluene, dichloromethane or tetrahydrofuran.

Compounds of formula (VII) are commercially available or they are known compounds or they are prepared by processes known in the art.

Process f): Compounds of formula (II) and compounds of formula (VIII) may be reacted together in the presence of a base, for example sodium t-butoxide, optionally in the presence of a catalyst for example tris(dibenzylidene-acetone)dipalladium(0) and S-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (0.113 g, 0.18 mM) in a suitable solvent such as toluene.

Compounds of formula (VIII) are commercially available or they are known compounds or they are prepared by processes known in the art.

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halo group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl. The reader is referred to Advanced Organic Chemistry, 4th Edition, by Jerry March, published by John Wiley & Sons 1992, for general guidance on reaction conditions and reagents.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard
5 practice (for illustration see T.W. Green, Protective Groups in Organic Synthesis, John Wiley and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a
10 methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali
15 metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with
20 a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an
25 arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by
30 hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis

- 32 -

with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

5 The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

Biological Assays

The activity of compounds of the invention was measured in a neuropeptide Y5 receptor binding assay as follows. Compounds were also tested in binding assays for the
10 neuropeptide Y₁ and neuropeptide Y₂ receptors. Activity against these 2 receptors is contraindicated for a neuropeptide Y5 antagonist.

a) Expression of human neuropeptide Y5 receptor in High 5TM insect cells.

High 5TM insect cells were obtained from Invitrogen (catalogue N° B855-02) and stored in liquid nitrogen. Cells were revived from liquid nitrogen storage and grown at 28°C
15 in 100 ml ExCell 405 (JRH Biosciences) serum free medium in a 250 ml conical flask (Corning) agitated at 140 rpm in an Innova 4330 orbital shaker (New Brunswick Scientific). Cultures were routinely sub-cultured every 3 - 4 days.

High 5TM insect cells were transfected with the human NPY5 receptor as follows. PCR primers were designed against the huNPY5 receptor sequence, Genbank Accession Number
20 U56079 [Gerald *et. al.* (1996) Nature 382, 168-171], but starting at base 56 through to base 1393, to express the protein 10 amino acid residues shorter at the amino terminal end [see Borowsky *et. al.* (1998) Regulatory Peptides 75-76, 45-53]. These primers were used to amplify the huNPY5 receptor from human placenta genomic DNA by PCR. This was then sub-cloned into pZERO2 (obtained from Invitrogen) for sequencing and re-cloned into
25 pFASTBAC1 (obtained from GIBCO BRL Life Technologies) for expression. Human NPY_r was isolated from pZERO2 on BamHI fragment and sub-cloned into pFastbac1 on BamHI restriction site. The junctions were sequenced to ensure correct prior to expression.

A baculovirus containing the pFASTBAC1 was then generated using the Bac-to-BacTM baculovirus expression system [Anderson *et al* (1996) FASEB Journal 10(6), 727-726]
30 (obtained from GIBCO BRL Life Technologies) following the protocol supplied with this expression system by GIBCO BRL Life Technologies.

- 33 -

High 5TM insect cells were infected with the baculovirus to transfect the cells with the human neuropeptide Y5 receptor as follows: Batches were grown for membrane preparation by inoculating 5 L of ExCell 405TM medium in a 7 L Bioreactor (FT-Applikon) with 1.75×10^9 mid log High 5TM cells. After 2-3 days growth at 28°C the mid log culture was infected with Baculovirus expressing the human NPY5 receptor at a multiplicity of infection (MOI) of 1.0. Cells (typically 1×10^{10}) were harvested 48 hours post infection by centrifugation (Heraeus Omnifuge 2.0RS 30 min, 296 g, 4°C) and flash frozen in liquid nitrogen for storage at -80°C.

b) Membrane preparation procedure

The following buffer was prepared daily and stored at 4°C. 50 mM Tris HCl pH 7.4, 5 mM EDTA and 10% w.v. sucrose. A protease inhibitor cocktail (Boehringer Mannheim) was added to both buffers according to the manufacturers instruction. Cells were thawed rapidly in three times their packed cell volume of hypotonic buffer (3:1 mix of water and buffer) and lysed routinely on ice using five Vibra Cell Sonicator (Sonics and Materials Inc.) bursts of ten seconds for the High 5TM insect cells. The cell lysate (typically 10-15 ml) was carefully loaded onto a 10 ml 41% sucrose cushion which was topped off with lysis buffer and spun at 150,000 g for 1 hour at 4°C in a Beckman Optima LE-80K Ultracentrifuge. The membrane fraction was carefully removed from the inter-phase and diluted at least four fold with lysis buffer. The membrane pellets were recovered by centrifugation at 150,000 g for 20 min at 4°C in a Beckman Optima LE-80K Ultracentrifuge and re-suspended at 5×10^7 cell equivalents per ml. The re-suspended membranes were divided into working aliquots, routinely 1 ml, flash frozen in liquid nitrogen and stored frozen at -80°C until use.

Prior to use the 1 ml High 5TM membranes were thawed and resuspended in 8 ml binding buffer (see below). Membranes are used at approximately 7 µg/ml of protein per incubate.

c) Neuropeptide Y5 receptor binding assay

The following reagents were used:

Binding buffer: 50 mM HEPES, 2.5 mM CaCl₂, 1 mM MgCl₂, 0.5% BSA, pH=7.4

Binding wash buffer: 50 mM HEPES, 2.5 mM CaCl₂, 1 mM MgCl₂, 0.5 M NaCl, 0.5% BSA, pH=7.4

Unifilter GFC filter plates: 50 µl of 0.5% polyethyleneimine was added to each well and left to equilibrate for four hours before use

Incubation plates: 96 well polypropylene plates, siliconised prior to use

Test Compounds: Compounds were dissolved in DMSO at a concentration of 1 mM. Final concentration of DMSO in the assay did not exceed 1%.

Peptide PYY (pancreatic polypeptide Y) - 10 μ M stock solution in binding buffer.

¹²⁵I PYY - 10 μ Ci/ml stock solution, diluted 1:10 dilution, into binding buffer.

5 Assays were performed in 96 well microtitre plates. 10 μ l of diluted test compound was added to each well of a plate, followed by 80 μ l of membranes and 10 μ l of radiolabelled ¹²⁵I PYY (0.01 μ Ci per well). Total and non-specific binding controls were included in each plate. The non-specific binding wells received 10 μ l of Peptide PYY from the 10 μ M stock solution, whilst the total binding wells received 10 μ l of binding buffer. For each assay, a
10 duplicate dose response of peptide PYY was included, top concentration 1 μ M.

The plates were incubated for two hours at room temperature with mixing, and then filtered onto the pre-treated filter plates. The incubation plates were washed twice with 150 μ l of cold binding wash buffer per well, then the filter plates were further washed with approximately 2.5 ml per well. The filter plates were dried overnight at room temperature, the
15 bottoms were sealed, and 20 μ l of Scintillant (Microscint 40, Canberra Packard) was added to each well. The tops of the plates were sealed and the plates were counted for 1 minute on a protocol set up for ¹²⁵I on a 96 well plate liquid scintillation counter (Top Count, Canberra Packard).

Compounds were considered to be active if they inhibited the binding by more than
20 50% at a concentration of 10 μ M. Dose responses were carried out on all compounds found to be active (8 point curves in duplicate).

Although the pharmacological properties of the compounds of the formula (I) vary with structural change as expected, in general compounds of the formula (I) possess an IC₅₀ in the above test in the range, for example, 0.0002 to 200 μ M. For example, the compound of
25 Example 16 has an IC₅₀ for the neuropeptide Y5 receptor of 94nM.

In order to use a compound of the formula (I) or a pharmaceutically acceptable salt, prodrug or solvate thereof, for the therapeutic treatment (including prophylactic treatment) of mammals including humans, it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

30 Therefore according to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the formula (I) or a

pharmaceutically acceptable salt, prodrug or solvate thereof, as defined herein before in association with a pharmaceutically acceptable diluent or carrier.

According to another aspect of the invention there is provided a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt, 5 prodrug or solvate thereof, in association with a pharmaceutically acceptable diluent or carrier for the treatment of disorders mediated by the neuropeptide Y5 receptor in a warm-blooded animal, such as a human being in need of such treatment.

To treat disorders mediated by the neuropeptide Y5 receptor neuropeptide Y5 receptor agonists or antagonists can be administered.

10 According to an additional aspect of the invention there is provided a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt, prodrug or solvate thereof, in association with a pharmaceutically acceptable diluent or carrier for the treatment of eating disorders in a warm-blooded animal, such as a human being.

15 According to an additional aspect of the invention there is provided a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt, prodrug or solvate thereof, in association with a pharmaceutically acceptable diluent or carrier for promoting weight loss in a warm-blooded animal, such as a human being.

20 Examples of disorders mediated by the neuropeptide Y5 receptor are eating disorders. Examples of eating disorders include obesity, bulimia or anorexia. Examples of eating disorders include obesity and related disorders, bulimia or anorexia. Examples of "related disorders" are diabetes dyslipidaemia, hypertension and sleep disturbances. Preferably "related disorders" refers to diabetes.

25 The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for 30 example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

5 Suitable pharmaceutically-acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or algenic acid; binding agents such as starch; lubricating agents such as magnesium stearate, stearic acid or talc; preservative agents such as ethyl or propyl *p*-hydroxybenzoate, and anti-oxidants, such as
10 ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the gastrointestinal tract, or to improve their stability and/or appearance, in either case, using conventional coating agents and procedures well known in the art.

 Compositions for oral use may be in the form of hard gelatin capsules in which the
15 active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin, or olive oil.

 Aqueous suspensions generally contain the active ingredient in finely powdered form together with one or more suspending agents, such as sodium carboxymethylcellulose,
20 methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation products of an alkylene oxide with fatty acids (for example polyoxethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters
25 derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl *p*-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening
30 agents (such as sucrose, saccharine or aspartame).

 Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid

paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

5 Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients such as sweetening, flavouring and colouring agents, may also be present.

10 The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, or a mineral oil, such as for example liquid paraffin or a mixture of any of these. Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, an esters or partial
15 esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring and preservative agents.

 Syrups and elixirs may be formulated with sweetening agents such as glycerol,
20 propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

 The pharmaceutical compositions may also be in the form of a sterile injectable aqueous or oily suspension, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which
25 have been mentioned above. A sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol.

 Suppository formulations may be prepared by mixing the active ingredient with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal
30 temperature and will therefore melt in the rectum to release the drug. Suitable excipients include, for example, cocoa butter and polyethylene glycols.

 Topical formulations, such as creams, ointments, gels and aqueous or oily solutions or

suspensions, may generally be obtained by formulating an active ingredient with a conventional, topically acceptable, vehicle or diluent using conventional procedures well known in the art.

Compositions for administration by insufflation may be in the form of a finely divided powder containing particles of average diameter of, for example, 30 μ m or much less, the powder itself comprising either active ingredient alone or diluted with one or more physiologically acceptable carriers such as lactose. The powder for insufflation is then conveniently retained in a capsule containing, for example, 1 to 50 mg of active ingredient for use with a turbo-inhaler device, such as is used for insufflation of the known agent sodium cromoglycate.

Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

For further information on formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient. For further information on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The size of the dose for therapeutic or prophylactic purposes of a compound of the formula (I) will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well known principles of medicine.


In using a compound of the formula (I) for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for example, 0.5 mg to 75 mg per kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous administration, a dose in the range, for example, 0.5 mg to 30 mg per kg body weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.5 mg to 25 mg per kg body weight will be used. Oral administration is however preferred, particularly in tablet form. Typically, unit dosage forms will contain about 1 mg to 500 mg of a compound of this invention.

The compounds of this invention may be used in combination with other drugs and therapies used in the treatment of disease states which would benefit from antagonism at the neuropeptide Y5 receptor. For example, the compounds of the formula (I) could be used in combination with drugs and therapies used in the treatment of eating disorders.

If formulated as a fixed dose such combination products employ the compounds of this invention within the dosage range described herein and the other pharmaceutically-active agent within its approved dosage range. Sequential use is contemplated when a combination formulation is inappropriate.

Although the compounds of the formula (I) are primarily of value as therapeutic agents for use in a warm-blooded animal, such as a human being, they are also useful whenever it is required to antagonise binding at the neuropeptide Y5 receptor. Thus, they are useful as pharmacological standards for use in the development of new biological tests and in the search for new pharmacological agents.

According to an additional aspect of the invention there is provided a compound of formula (IA):



(IA)

wherein:

R^1 is cyano, halo, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl, C_{1-4} alkoxy, N -(C_{1-4} alkyl)amino or N,N -(C_{1-4} alkyl) $_2$ amino;

R^2 is halo, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl or C_{1-4} alkoxy;

A is $—NR^a—$, $—O—$ or a direct bond; wherein R^a is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl; wherein said C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl may be optionally substituted by one or more R^4 ;

5 R^3 is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl or C_{2-10} alkynyl wherein said C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl may be optionally substituted by one or more R^5 ; or R^3 is carbocyclyl or heterocyclyl wherein said carbocyclyl or heterocyclyl may be optionally substituted on carbon by one or more R^6 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R^7 ;

10 R^4 and R^5 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, C_{1-6} alkanoylamino, C_{2-6} alkenyloxy, carbonyl, C_{1-6} alkoxycarbonyl, heterocyclyl, heterocycliloxy, heterocyclylcarbonyl, heterocycliloxy, carbocyclyl, carbocycliloxy, carbocyclylcarbonyl, carbocycliloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkoxycarbonylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkylS(O)_a wherein a is 0-2, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl)₂sulphamoyl; wherein any C_{1-6} alkyl, heterocyclyl or carbocyclyl may be optionally substituted on carbon by one or more R^6 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R^7 ;

20 R^6 is selected from halo, hydroxy, cyano, carbamoyl, ureido, trifluoromethyl, trifluoromethoxy, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, C_{2-4} alkenyloxy, carbonyl, C_{1-4} alkanoyl, C_{1-4} alkanoylamino, C_{1-4} alkanoyloxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0-2, N -(C_{1-4} alkyl)sulphamoyl, N,N -(C_{1-4} alkyl)₂sulphamoyl, heterocyclyl, heterocycliloxy, heterocyclylcarbonyl, heterocycliloxy, carbocyclyl, carbocycliloxy, carbocyclylcarbonyl and carbocycliloxy; wherein any C_{1-4} alkyl, carbocyclyl and heterocyclyl may be optionally substituted on carbon by one or more R^8 ;

R^7 is selected from C_{1-4} alkyl, C_{1-4} alkanoyl, C_{1-4} alkylsulphonyl, C_{1-4} alkoxycarbonyl, carbamoyl, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, benzyl, benzoyl, phenylsulphonyl and phenyl;

30

R^8 is selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, methoxy, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, *N*-methylcarbamoyl, *N,N*-dimethylcarbamoyl, methylthio, methylsulphanyl, mesyl, *N*-methylsulphamoyl, *N,N*-dimethylsulphamoyl, heterocyclyl, heterocyclyloxy, heterocyclylcarbonyl, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylcarbonyl and carbocyclyloxycarbonyl;

x is 0 - 4; wherein the values of R^1 may be the same or different;

y is 0 - 3; wherein the values of R^2 may be the same or different; and

10 n is 0 - 2;

or a pharmaceutically acceptable salt, prodrug or solvate thereof;

with the provisos:

1) when R^3 is a nitrogen linked heterocyclyl, A is a direct bond;

2) when x and y are 0 and n is 2, the group R^3 -A-C(O)-NH- is not 2-formamido, 2-acetamido, 15 2-propionamido, 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido), 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido), 2-(2-phthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido), 2-(2-amino-2-benzylacetamido), 3-(2-phthalimidoacetamido), 3-(3-phthalimidopropionamido), 20 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido), 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido), 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-benzylacetamido) or 2-(3,3,3-trifluoro-2-hydroxy-2-methylpropionamido);


3) when x and y are 0 and n is 1, the group R^3 -A-C(O)-NH- is not 3-(2-phthalimidoacetamido), 25 3-(3-phthalimidopropionamido), 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido), 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido), 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-isobutylacetamido) or 3-(2-amino-2-benzylacetamido);

4) when x and y are 0 and n is 0, the group R^3 -A-C(O)-NH- is not 2-(2-phthalimidoacetamido), 30 2-(3-phthalimidopropionamido), 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido), 2-(2-phthalimido-2-benzylacetamido),

2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido),
2-(2-amino-2-isobutylacetamido) or 2-(2-amino-2-benzylacetamido);
5) when $(R^1)_x$ is 7-fluoro, y is 0 and n is 2, the group $R^3-A-C(O)-NH-$ is not 3-acetamido;
or a pharmaceutically acceptable salt, prodrug or solvate thereof, for use as a medicament.



5 In particular additional aspect of the invention there is provided a compound of formula (IA) which is 2-acetamidodibenzothiophene or a pharmaceutically acceptable salt, prodrug or solvate thereof, for use as a medicament.

According to an additional aspect of the invention there is provided a compound of formula (IA'):



wherein:

X is a group of formula (A) or (B):



15

R^1 is cyano, halo, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl, C_{1-4} alkoxy, $N-(C_{1-4}alkyl)amino$ or $N,N-(C_{1-4}alkyl)_2amino$;

R^2 is halo, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl or C_{1-4} alkoxy;

A is $-NR^a-$, $-O-$ or a direct bond; wherein R^a is hydrogen, C_{1-10} alkyl,

20 $C_{2-10}alkenyl$, $C_{2-10}alkynyl$; wherein R^a may be optionally substituted by one or more R^5 ;

R^3 is hydrogen, C_{1-10} alkyl, $C_{2-10}alkenyl$ or $C_{2-10}alkynyl$ wherein R^3 may be optionally substituted by one or more R^6 ; or R^3 is carbocyclyl or heterocyclyl wherein R^3 may be optionally substituted on carbon by one or more R^7 ; and wherein if said heterocyclyl contains an $-NH-$ moiety that nitrogen may be optionally substituted by R^8 ;

25 R^4 is halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, $C_{1-4}alkoxycarbonyl$, $C_{1-4}alkyl$, $C_{2-4}alkenyl$, $C_{2-4}alkynyl$, $C_{1-4}alkoxy$, $C_{1-4}alkanoyl$, $C_{1-4}alkanoyloxy$, $N-(C_{1-4}alkyl)amino$, $N,N-(C_{1-4}alkyl)_2amino$,

- C_{1-4} alkanoylamino, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl,
 N -(carbocyclyl)carbamoyl, N,N -(carbocyclyl)₂carbamoyl, N -(heterocyclyl)carbamoyl,
 N,N -(heterocyclyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0 to 2, C_{1-4} alkoxycarbonyl,
 N -(C_{1-4} alkyl)sulphamoyl, N,N -(C_{1-4} alkyl)₂sulphamoyl, C_{1-4} alkylsulphonylamino or
 5 (nitrogen-linked heterocyclic ring)carbonyl;
- R^5 and R^6 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido,
 amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, C_{1-6} alkoxy,
 C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, C_{1-6} alkanoylamino, C_{2-6} alkenyloxy, carbonyl,
 C_{1-6} alkoxycarbonyl, heterocyclyl, heterocycliloxy, heterocyclylamino,
 10 heterocyclyl- N -(C_{1-6} alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl,
 heterocyclylcarbonylamino, heterocycliloxy, carbocyclyl, carbocycliloxy,
 carbocyclylamino, carbocyclyl- N -(C_{1-6} alkyl)amino, carbocyclylsulphonyl,
 carbocyclylcarbonyl, carbocyclylcarbonylamino, carbocycliloxy, N -(C_{1-6} alkyl)amino,
 N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkoxycarbonylamino, N -(C_{1-6} alkyl)carbamoyl,
 15 N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkylS(O)_a wherein a is 0-2, N -(C_{1-6} alkyl)sulphamoylamino,
 N,N -(C_{1-6} alkyl)₂sulphamoylamino, C_{1-6} alkylsulphonylamino,
 (C_{1-6} alkyl)sulphonyl- N -(C_{1-6} alkyl)amino, N -(C_{1-6} alkyl)sulphamoyl and
 N,N -(C_{1-6} alkyl)₂sulphamoyl; wherein R^5 and R^6 may be optionally substituted on carbon by
 one or more R^9 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may
 20 be independently optionally substituted by R^{10} ;
- R^7 and R^9 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido,
 trifluoromethyl, trifluoromethoxy, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl,
 aminosulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl,
 C_{1-4} alkoxycarbonylamino, C_{2-4} alkenyloxy, carbonyl, C_{1-4} alkanoyl, C_{1-4} alkanoylamino,
 25 C_{1-4} alkanoyloxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, N -(C_{1-4} alkyl)carbamoyl,
 N,N -(C_{1-4} alkyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0-2, N -(C_{1-4} alkyl)sulphamoylamino,
 N,N -(C_{1-4} alkyl)₂sulphamoylamino, (C_{1-4} alkyl)sulphonylamino,
 (C_{1-4} alkyl)sulphonyl- N -(C_{1-4} alkyl)amino, N -(C_{1-4} alkyl)sulphamoyl,
 N,N -(C_{1-4} alkyl)₂sulphamoyl, heterocyclyl, heterocycliloxy, heterocyclylamino,
 30 heterocyclyl- N -(C_{1-4} alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl,
 heterocyclylcarbonylamino, heterocycliloxy, carbocyclyl, carbocycliloxy,
 carbocyclylamino, carbocyclyl- N -(C_{1-4} alkyl)amino, carbocyclylsulphonyl,

carbocyclylcarbonyl, carbocyclylcarbonylamino, and carbocyclyloxycarbonyl; wherein R^7 and R^9 may be independently optionally substituted on carbon by one or more R^{11} ;

R^8 and R^{10} are independently selected from C_{1-4} alkyl, C_{1-4} alkanoyl, sulphamoyl, C_{1-4} alkylsulphonyl, C_{1-4} alkoxycarbonyl, carbamoyl, N -(C_{1-4} alkyl)carbamoyl, 5 N,N -(C_{1-4} alkyl)₂carbamoyl, N -(C_{1-4} alkyl)sulphamoyl, N,N -(C_{1-4} alkyl)₂sulphamoyl, heterocyclyl, heterocyclylcarbonyl, heterocyclyloxycarbonyl, heterocyclylsulphonyl, carbocyclyl, carbocyclylcarbonyl, carbocyclyloxycarbonyl and carbocyclylsulphonyl; wherein R^8 and R^{10} may be independently optionally substituted on carbon by one or more R^{12} ;

R^{11} and R^{12} are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, 10 amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, vinyl, allyl, methoxy, ethoxy, vinyloxy, allyloxy, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, N -methylcarbamoyl, N,N -dimethylcarbamoyl, methylthio, methylsulphanyl, mesyl, N -methylsulphamoyl, N,N -dimethylsulphamoyl, heterocyclyl, heterocyclyloxy, heterocyclylcarbonyl, heterocyclyloxycarbonyl, carbocyclyl, 15 carbocyclyloxy, carbocyclylcarbonyl and carbocyclyloxycarbonyl;

x is 0 - 4; wherein the values of R^1 may be the same or different;

y is 0 - 3; wherein the values of R^2 may be the same or different;

z is 0 - 3; wherein the values of R^4 may be the same or different; and

n is 0 - 2;


20 or a pharmaceutically acceptable salt, prodrug or solvate thereof;

with the provisos when X is a group of formula (A):

- 1) when R^3 is a nitrogen linked heterocyclyl, A is a direct bond;
- 2) when x and y are 0 and n is 2, the group R^3 -A-C(O)-NH- is not 2-formamido, 2-acetamido, 2-propionamido, 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido), 25 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido), 2-(2-phthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido), 2-(2-amino-2-benzylacetamido), 3-(2-phthalimidoacetamido), 3-(3-phthalimidopropionamido), 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido), 30 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido), 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-benzylacetamido) or 2-(3,3,3-trifluoro-2-hydroxy-2-methylpropionamido);

- 3) when x and y are 0 and n is 1, the group R^3 -A-C(O)-NH- is not 3-(2-phthalimidoacetamido), 3-(3-phthalimidopropionamido), 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido), 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido), 3-(2-amino-2-isopropylacetamido), 5 3-(2-amino-2-isobutylacetamido) or 3-(2-amino-2-benzylacetamido);
- 4) when x and y are 0 and n is 0, the group R^3 -A-C(O)-NH- is not 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido), 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido), 2-(2-phthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido), 10 2-(2-amino-2-isobutylacetamido) or 2-(2-amino-2-benzylacetamido);
- 5) when $(R^1)_x$ is 7-fluoro, y is 0 and n is 2, the group R^3 -A-C(O)-NH- is not 3-acetamido; or a pharmaceutically acceptable salt, prodrug or solvate thereof, for use as a medicament.

- According to a further feature of the invention there is provided a method of treating disorders mediated by the neuropeptide Y5 receptor in a warm-blooded animal, such as a 15 human being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB):



(IB)

wherein:

- 20 R^1 is cyano, halo, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl, C_{1-4} alkoxy, N -(C_{1-4} alkyl)amino or N,N -(C_{1-4} alkyl)₂amino;
- R^2 is halo, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl or C_{1-4} alkoxy;
- A is $-NR^a$, $-O-$ or a direct bond; wherein R^a is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl; wherein said C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl may be optionally 25 substituted by one or more R^4 ;
- R^3 is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl or C_{2-10} alkynyl wherein said C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl may be optionally substituted by one or more R^5 ; or R^3 is carbocyclyl or heterocyclyl wherein said carbocyclyl or heterocyclyl may be optionally

substituted on carbon by one or more R⁶; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R⁷;

- R⁴ and R⁵ are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, C₁₋₆alkanoylamino, C₂₋₆alkenyloxy, carbonyl, C₁₋₆alkoxycarbonyl, heterocyclyl, heterocycliloxy, heterocyclylcarbonyl, heterocycliloxy, carbocyclyl, carbocycliloxy, carbocyclylcarbonyl, carbocycliloxy, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkoxycarbonylamino, N-(C₁₋₆alkyl)carbamoyl, N,N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkylS(O)_a wherein a is 0-2, N-(C₁₋₆alkyl)sulphamoyl and N,N-(C₁₋₆alkyl)₂sulphamoyl; wherein any C₁₋₆alkyl, heterocyclyl or carbocyclyl may be optionally substituted on carbon by one or more R⁶; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R⁷;

- R⁶ is selected from halo, hydroxy, cyano, carbamoyl, ureido, trifluoromethyl, trifluoromethoxy, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy, C₁₋₄alkoxycarbonyl, C₂₋₄alkenyloxy, carbonyl, C₁₋₄alkanoyl, C₁₋₄alkanoylamino, C₁₋₄alkanoyloxy, N-(C₁₋₄alkyl)amino, N,N-(C₁₋₄alkyl)₂amino, N-(C₁₋₄alkyl)carbamoyl, N,N-(C₁₋₄alkyl)₂carbamoyl, C₁₋₄alkylS(O)_a wherein a is 0-2, N-(C₁₋₄alkyl)sulphamoyl, N,N-(C₁₋₄alkyl)₂sulphamoyl, heterocyclyl, heterocycliloxy, heterocyclylcarbonyl, heterocycliloxy, carbocyclyl, carbocycliloxy, carbocyclylcarbonyl and carbocycliloxy; wherein any C₁₋₄alkyl, carbocyclyl and heterocyclyl may be optionally substituted on carbon by one or more R⁸;

R⁷ is selected from C₁₋₄alkyl, C₁₋₄alkanoyl, C₁₋₄alkylsulphonyl, C₁₋₄alkoxycarbonyl, carbamoyl, N-(C₁₋₄alkyl)carbamoyl, N,N-(C₁₋₄alkyl)₂carbamoyl, benzyl, benzoyl, phenylsulphonyl and phenyl;

- R⁸ is selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, methoxy, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, N-methylcarbamoyl, N,N-dimethylcarbamoyl, methylthio, methylsulphonyl, mesyl, N-methylsulphamoyl, N,N-dimethylsulphamoyl, heterocyclyl, heterocycliloxy, heterocyclylcarbonyl, heterocycliloxy, carbocyclyl, carbocycliloxy, carbocyclylcarbonyl and carbocycliloxy;

x is 0 - 4; wherein the values of R¹ may be the same or different;

- 47 -

y is 0 - 3; wherein the values of R^2 may be the same or different; and

n is 0 - 2;


or a pharmaceutically acceptable salt, prodrug or solvate thereof;

with the providing that when R^3 is a nitrogen linked heterocyclyl, A is a direct bond;

5 or a pharmaceutically acceptable salt, prodrug or solvate thereof.

According to a particular feature of the invention there is provided a method for promoting weight loss in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB) or a pharmaceutically acceptable salt, prodrug or solvate thereof.

10 According to a further feature of the invention there is provided a method of treating disorders mediated by the neuropeptide Y5 receptor in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB'):




(IB')


15

wherein:

X is a group of formula (A) or (B):



(A)



(B)

20 R^1 is cyano, halo, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl, C_{1-4} alkoxy, N -(C_{1-4} alkyl)amino or N,N -(C_{1-4} alkyl)₂amino;

R^2 is halo, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl or C_{1-4} alkoxy;

A is $-NR^a-$, $-O-$ or a direct bond; wherein R^a is hydrogen, C_{1-10} alkyl,

C_{2-10} alkenyl, C_{2-10} alkynyl; wherein R^a may be optionally substituted by one or more R^5 ;

25 R^3 is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl or C_{2-10} alkynyl wherein R^3 may be optionally substituted by one or more R^6 ; or R^3 is carbocyclyl or heterocyclyl wherein R^3 may be

optionally substituted on carbon by one or more R^7 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R^8 ;

- R^4 is halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-4} alkoxycarbonyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl,
- 5 C_{1-4} alkoxy, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, C_{1-4} alkanoylamino, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, N -(heterocyclyl)carbamoyl, N,N -(heterocyclyl)₂carbamoyl, N -(heterocyclyl)carbamoyl, N,N -(heterocyclyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0 to 2, C_{1-4} alkoxycarbonyl, N -(C_{1-4} alkyl)sulphamoyl, N,N -(C_{1-4} alkyl)₂sulphamoyl, C_{1-4} alkylsulphonylamino or
- 10 (nitrogen-linked heterocyclic ring)carbonyl;
- R^5 and R^6 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, C_{1-6} alkanoylamino, C_{2-6} alkenyloxycarbonyl, C_{1-6} alkoxycarbonyl, heterocyclyl, heterocycliloxy, heterocyclylamino,
- 15 heterocyclyl- N -(C_{1-6} alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl, heterocyclylcarbonylamino, heterocycliloxycarbonyl, carbocyclyl, carbocycliloxy, carbocyclylamino, carbocyclyl- N -(C_{1-6} alkyl)amino, carbocyclylsulphonyl, carbocyclylcarbonyl, carbocyclylcarbonylamino, carbocycliloxycarbonyl, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkoxycarbonylamino, N -(C_{1-6} alkyl)carbamoyl,
- 20 N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkylS(O)_a wherein a is 0-2, N -(C_{1-6} alkyl)sulphamoylamino, N,N -(C_{1-6} alkyl)₂sulphamoylamino, C_{1-6} alkylsulphonylamino, (C_{1-6} alkyl)sulphonyl- N -(C_{1-6} alkyl)amino, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl)₂sulphamoyl; wherein R^5 and R^6 may be optionally substituted on carbon by one or more R^9 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may
- 25 be independently optionally substituted by R^{10} ;

- R^7 and R^9 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, trifluoromethyl, trifluoromethoxy, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, C_{1-4} alkoxycarbonylamino, C_{2-4} alkenyloxycarbonyl, C_{1-4} alkanoyl, C_{1-4} alkanoylamino,
- 30 C_{1-4} alkanoyloxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0-2, N -(C_{1-4} alkyl)sulphamoylamino, N,N -(C_{1-4} alkyl)₂sulphamoylamino, (C_{1-4} alkyl)sulphonylamino,

(C₁₋₄alkyl)sulphonyl-*N*-(C₁₋₄alkyl)amino, *N*-(C₁₋₄alkyl)sulphamoyl, *N,N*-(C₁₋₄alkyl)₂sulphamoyl, heterocyclyl, heterocycloxy, heterocyclylamino, heterocyclyl-*N*-(C₁₋₄alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl, heterocyclylcarbonylamino, heterocycloxy carbonyl, carbocyclyl, carbocycloxy, carbocyclylamino, carbocyclyl-*N*-(C₁₋₄alkyl)amino, carbocyclylsulphonyl, carbocyclylcarbonyl, carbocyclylcarbonylamino, and carbocycloxy carbonyl; wherein R⁷ and R⁹ may be independently optionally substituted on carbon by one or more R¹¹;

R⁸ and R¹⁰ are independently selected from C₁₋₄alkyl, C₁₋₄alkanoyl, sulphamoyl, C₁₋₄alkylsulphonyl, C₁₋₄alkoxycarbonyl, carbamoyl, *N*-(C₁₋₄alkyl)carbamoyl, *N,N*-(C₁₋₄alkyl)₂carbamoyl, *N*-(C₁₋₄alkyl)sulphamoyl, *N,N*-(C₁₋₄alkyl)₂sulphamoyl, heterocyclyl, heterocyclylcarbonyl, heterocycloxy carbonyl, heterocyclylsulphonyl, carbocyclyl, carbocyclylcarbonyl, carbocycloxy carbonyl and carbocyclylsulphonyl; wherein R⁸ and R¹⁰ may be independently optionally substituted on carbon by one or more R¹²;

R¹¹ and R¹² are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, vinyl, allyl, methoxy, ethoxy, vinyloxy, allyloxy, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, *N*-methylcarbamoyl, *N,N*-dimethylcarbamoyl, methylthio, methylsulphinyl, mesyl, *N*-methylsulphamoyl, *N,N*-dimethylsulphamoyl, heterocyclyl, heterocycloxy, heterocyclylcarbonyl, heterocycloxy carbonyl, carbocyclyl, carbocycloxy, carbocyclylcarbonyl and carbocycloxy carbonyl;

x is 0 - 4; wherein the values of R¹ may be the same or different;

y is 0 - 3; wherein the values of R² may be the same or different;

z is 0 - 3; wherein the values of R⁴ may be the same or different; and

n is 0 - 2;

with the providing that when R³ is a nitrogen linked heterocyclyl, A is a direct bond; or a pharmaceutically acceptable salt, prodrug or solvate thereof.

According to a particular feature of the invention there is provided a method of treating eating disorders in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB') or a pharmaceutically acceptable salt, prodrug or solvate thereof.

According to a particular feature of the invention there is provided a method for promoting weight loss in a warm-blooded animal, such as a human being, in need of such

treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB') or a pharmaceutically acceptable salt, prodrug or solvate thereof.

In particular further feature of the invention there is provided a method of treating disorders mediated by the neuropeptide Y5 receptor in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB) which is 5,5-dioxo-2-[(2,2,2-trifluoro-1-hydroxy-1-methylethyl)carbonylamino]dibenzothiophene, 2-acetamidodibenzothiophene or 5,5-dioxo-2-acetamidodibenzothiophene or a pharmaceutically acceptable salt, prodrug or solvate thereof.

10 In particular further feature of the invention there is provided a method of treating eating disorders in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB') which is 5,5-dioxo-2-[(2,2,2-trifluoro-1-hydroxy-1-methylethyl)carbonylamino]dibenzothiophene, 15 2-acetamidodibenzothiophene or 5,5-dioxo-2-acetamidodibenzothiophene or a pharmaceutically acceptable salt, prodrug or solvate thereof.

According to this further feature of the invention there is provided a method of treating eating disorders in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a 20 compound of formula (IB) or a pharmaceutically acceptable salt, prodrug or solvate thereof.

According to this further feature of the invention there is provided a method of treating eating disorders in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB) which is 25 5,5-dioxo-2-[(2,2,2-trifluoro-1-hydroxy-1-methylethyl)carbonylamino]dibenzothiophene, 2-acetamidodibenzothiophene or 5,5-dioxo-2-acetamidodibenzothiophene or a pharmaceutically acceptable salt, prodrug or solvate thereof.

According to an additional aspect of the invention there is provided the use of a compound of formula (IB) or a pharmaceutically acceptable salt, prodrug or solvate thereof in the manufacture of a medicament for the treatment of disorders mediated by the 30 neuropeptide Y5 receptor in a warm-blooded animal, such as a human being.

- 51 -

According to an additional aspect of the invention there is provided the use of a compound of formula (IB') or a pharmaceutically acceptable salt, prodrug or solvate thereof in the manufacture of a medicament for the treatment of disorders mediated by the neuropeptide Y5 receptor in a warm-blooded animal, such as a human being.

5 According to an additional aspect of the invention there is provided the use of a compound of formula (IB) which is
5,5-dioxo-2-[(2,2,2-trifluoro-1-hydroxy-1-methylethyl)carbonylamino]dibenzothiophene,
2-acetamidodibenzothiophene or 5,5-dioxo-2-acetamidodibenzothiophene or a
pharmaceutically acceptable salt, prodrug or solvate thereof in the manufacture of a
10 medicament for the treatment of disorders mediated by the neuropeptide Y5 receptor in a warm-blooded animal, such as a human being.

 According to an additional aspect of the invention there is provided the use of a compound of formula (IB') which is
5,5-dioxo-2-[(2,2,2-trifluoro-1-hydroxy-1-methylethyl)carbonylamino]dibenzothiophene,
15 2-acetamidodibenzothiophene or 5,5-dioxo-2-acetamidodibenzothiophene or a pharmaceutically acceptable salt, prodrug or solvate thereof in the manufacture of a medicament for the treatment of eating disorders in a warm-blooded animal, such as a human being.

 According to an additional aspect of the invention there is provided the use of a
20 compound of formula (IB) or a pharmaceutically acceptable salt, prodrug or solvate thereof in the manufacture of a medicament for the treatment of eating disorders in a warm-blooded animal, such as a human being.

 According to an additional aspect of the invention there is provided the use of a compound of formula (IB') or a pharmaceutically acceptable salt, prodrug or solvate
25 thereof in the manufacture of a medicament for the treatment of eating disorders in a warm-blooded animal, such as a human being.

 According to an additional aspect of the invention there is provided the use of a compound of formula (IB) or a pharmaceutically acceptable salt, prodrug or solvate thereof in the manufacture of a medicament for promoting weight loss in a warm-blooded
30 animal, such as a human being.

 According to an additional aspect of the invention there is provided the use of a compound of formula (IB') or a pharmaceutically acceptable salt, prodrug or solvate

thereof in the manufacture of a medicament for promoting weight loss in a warm-blooded animal, such as a human being.

According to an additional aspect of the invention there is provided the use of a compound of formula (IB) which is

5 5,5-dioxo-2-[(2,2,2-trifluoro-1-hydroxy-1-methylethyl)carbonylamino]dibenzothiophene, 2-acetamidodibenzothiophene or 5,5-dioxo-2-acetamidodibenzothiophene or a pharmaceutically acceptable salt, prodrug or solvate thereof in the manufacture of a medicament for the treatment of eating disorders in a warm-blooded animal, such as a human being.

10 According to an additional aspect of the invention there is provided the use of a compound of formula (IB') which is 5,5-dioxo-2-[(2,2,2-trifluoro-1-hydroxy-1-methylethyl)carbonylamino]dibenzothiophene, 2-acetamidodibenzothiophene or 5,5-dioxo-2-acetamidodibenzothiophene or a pharmaceutically acceptable salt, prodrug or solvate thereof in the manufacture of a
15 medicament for the treatment of eating disorders in a warm-blooded animal, such as a human being.

According to another aspect of the invention there is provided a pharmaceutical composition comprising a compound of formula (IB'), or a pharmaceutically acceptable salt, prodrug or solvate thereof, in association with a pharmaceutically acceptable diluent
20 or carrier for the treatment of disorders mediated by the neuropeptide Y5 receptor in a warm-blooded animal, such as a human being.

According to another aspect of the invention there is provided a pharmaceutical composition comprising a compound of formula (IB'), or a pharmaceutically acceptable salt, prodrug or solvate thereof, in association with a pharmaceutically acceptable diluent
25 or carrier for the treatment of eating disorders in a warm-blooded animal, such as a human being.

According to another aspect of the invention there is provided a pharmaceutical composition comprising a compound of formula (IB'), or a pharmaceutically acceptable salt, prodrug or solvate thereof, in association with a pharmaceutically acceptable diluent
30 or carrier for promoting weight loss in a warm-blooded animal, such as a human being.

Examples

The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

- (i) temperatures are given in degrees Celsius ($^{\circ}\text{C}$); operations were carried out at room or
5 ambient temperature, that is, at a temperature in the range of $18\text{--}25^{\circ}\text{C}$;
- (ii) organic solutions were dried over anhydrous magnesium sulphate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure ($600\text{--}4000$ Pascals; $4.5\text{--}30$ mmHg) with a bath temperature of up to 60°C ;
- (iii) chromatography means flash chromatography on silica gel (Merck Keisegel ART 9385);
10 thin layer chromatography (TLC) was carried out on silica gel plates; where a "Bond Elut" column is referred to, this means a column containing 20 g of silica, the silica being contained in a 70 ml disposable syringe and supported by a porous disc of 54\AA pore size, obtained from International Sorbent Technology under the name "ISOLUTE"; "ISOLUTE" is a registered trade mark;
- 15 (iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;
- (v) final products had satisfactory proton nuclear magnetic resonance (NMR) spectra and/or mass spectral data;
- (vi) yields are given for illustration only and are not necessarily those which can be obtained
20 by diligent process development; preparations were repeated if more material was required;
- (vii) when given, NMR data is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard, determined at 300 MHz using perdeuterio dimethyl sulphoxide ($\text{DMSO-}d_6$) as solvent unless otherwise indicated; s, singlet; d, doublet; dd, double doublet; t, triplet; tt, triple triplet; q,
25 quartet; tq, triple quartet; m, multiplet; br, broad;
- (viii) chemical symbols have their usual meanings; SI units and symbols are used;
- (ix) solvent ratios are given in volume:volume (v/v) terms; and
- (x) mass spectra were run with an electron energy of 70 electron volts in the chemical
ionization (CI) mode using a direct exposure probe; where indicated ionization was effected
30 by electron impact (EI), fast atom bombardment (FAB) or electrospray (ES); values for m/z are given; generally, only ions which indicate the parent mass are reported and unless otherwise stated, the $(\text{MH})^+$ is quoted;

(xi) unless stated otherwise compounds containing an asymmetrically substituted carbon and/or sulphur atom have not been resolved;

(xii) where a synthesis is described as being analogous to that described in a previous example the amounts used are the millimolar ratio equivalents to those used in the previous example;

5 (xvi) the following abbreviations have been used:

	SM	starting material;
	EDAC	1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride;
	EtOAc	ethyl acetate;
	THF	tetrahydrofuran;
10	DCM	dichloromethane;
	DMA	dimethylacetamide;
	ether	diethyl ether;
	DMF	<i>N,N</i> -dimethylformamide; and
	DMSO	dimethylsulphoxide; and

15 xvii) where examples have chiral centres they are racemic mixtures unless otherwise indicated.

Example 1

2-(4-Nitrobenzoxycarbonylamino)dibenzothiophene

20 To a solution of *p*-nitrophenylchloroformate (5.57 g, 27.5 mmol) in EtOAc (60 ml) was added a solution of 2-aminodibenzthiophene (Bull. Soc. Chim. Fr. (1996), 133(6), 597-610; 5.0 g, 25 mmol) in EtOAc (100 ml) over a period of 30 minutes with ice bath cooling. The mixture was stirred for 1 hour then poured onto water (500 ml) and extracted with EtOAc (4 x 250 ml). The combined organic layers were washed with water (2 x 100 ml)
25 and brine (100 ml), dried and evaporated *in vacuo* to obtain the title compound (7.1 g). NMR: 10.66 (brs, 1H), 8.5 (s, 1H), 8.3 (d, 2H), 8.2 (m, 1H), 8.0 (m, 2H), 7.6 (m, 1H), 7.55 (d, 2H), 7.5 (m, 2H).

Example 2

30 2-(*t*-Butylcarbonylamino)dibenzothiophene

2-Aminodibenzthiophene (Bull. Soc. Chim. Fr. (1996), 133 (6), 597-610; 2.0 g, 10 mmol) and triethylamine (1.11 g, 11 mmol) were dissolved in DCM (50 ml) and cooled in an

ice bath. A solution of pivaloyl chloride (1.21 g, 10 mmol) in DCM (10 ml) was added slowly and the resultant solution was stirred at ambient temperature for 16 hours then evaporated *in vacuo*. The residue was triturated with methanol to leave the product as a colourless solid (1.85 g). NMR: 9.40 (s, 1H), 8.59 (s, 1H), 8.17 (m, 1H), 7.98 (d, 1H), 7.90 (d, 1H), 7.73 (dd, 5 1H), 7.46 (m, 2H), 1.15 (s, 9H); m/z 284.


Example 3

2-(1,2,4-Triazol-1-ylmethylcarbonylamino)dibenzothiophene












2-Aminodibenzthiophene (Bull. Soc. Chim. Fr. (1996), 133 (6), 597-610; 500 mg, 2.5 10 mmol) was dissolved DMF (10 ml) and 2-(1,2,4-triazol-1-yl)acetic acid (318 mg, 2.5 mmol) was added in one portion followed by 1-hydroxybenztriazole hydrate (383 mg, 5.0 mmol) and EDAC (960 mg, 5.0 mmol). The resultant mixture was stirred for 16 hours at ambient temperature then poured on to water (100 ml). A fine white solid was collected which was washed extensively with water, methanol (25 ml) and ether (25 ml) to leave a colourless solid 15 (650 mg). NMR: 10.62 (brs, 1H), 8.18 (m, 2H), 8.16 (m, 1H), 7.99 (m, 3H), 7.62 (dd, 1H), 7.50 (m, 2H), 5.20 (s, 2H); m/z 309.

Examples 4-18



Following the procedure of Example 3 using 2-aminodibenzthiophene (Bull. Soc. 20 Chim. Fr. (1996), 133 (6), 597-610) and the appropriate acid the following compounds were prepared.



Ex	R ¹	NMR	m/z
4		10.38 (brs, 1H), 8.6 (d, 1H), 8.15 (m, 1H), 8.0 (m, 1H), 7.9 (d, 1H), 7.6 (dd, 1H), 7.5 (m, 2H), 1.8 (m, 1H), 0.8 (m, 4H)	268
5		10.26 (brs, 1H), 8.59 (d, 1H), 8.17 (m, 1H), 8.00 (m, 1H), 7.94 (d, 1H), 7.61 (dd, 1H), 7.50 (m, 2H), 4.08 (s, 2H), 3.47 (t, 2H), 2.31 (t, 2H), 2.00 (m, 2H)	325

Ex	R ¹	m/z
6		284
7		360
8		327
9		312
10		348
11		347
12		323
13		335
14		320 (M-H) ⁻
15		375
16		355

- 57 -


17		285 (des CO ₂ C(CH ₃) ₃)
18		265 (M-H) ⁻


Example 192-(N'-Pyrid-4-ylethyl-N'-methylureido)dibenzothiophene



2-(4-Nitrobenzoxycarbonylamino)dibenzothiophene (Example 1; 501 mg, 1.35 mmol) was suspended in EtOAc (10 ml) and 4-[2-(methylamino)ethyl]pyridine (408 mg, 3.0 mmol) added in one portion. The resulting mixture was stirred at ambient temperature for 16 hours then further EtOAc (50 ml) added. The solution was washed with 1 M sodium hydroxide (3 x 25 ml), water (2 x 25 ml) and brine (25 ml), dried and evaporated *in vacuo*. NMR 8.44 (m, 3H), 8.34 (d, 1H), 8.13 (m, 1H), 7.96 (m, 1H), 7.82 (d, 1H), 7.55 (dd, 1H), 7.47 (m, 2H), 7.28 (d, 2H), 3.61 (t, 2H), 2.96 (s, 3H), 2.86 (t, 2H); m/z 362.

Examples 20-22

Following the procedure of Example 19 using 2-(4-nitrobenzoxycarbonylamino)dibenzothiophene (Example 1) and the appropriate amine the following compounds were prepared.



Ex	R ¹	NMR	m/z
20		8.72 (s, 1H), 8.40 (s, 1H), 8.14 (m, 1H), 7.96 (m, 1H), 7.84 (d, 1H), 7.58 (d, 1H), 7.46 (m, 2H), 3.63 (t, 4H), 3.45 (t, 4H)	313

21		8.64 (s, 1H), 8.43 (d, 1H), 8.13 (d, 2H), 7.97 (m, 1H), 7.85 (d, 1H), 7.62 (dd, 1H), 7.47 (m, 2H), 7.28 (d, 2H), 4.63 (s, 2H), 3.45 (q, 2H), 1.12 (t, 3H)	362
22		8.63 (s, 1H), 8.40 (d, 1H), 8.12 (m, 1H), 7.95 (m, 1H), 7.82 (d, 1H), 7.58 (dd, 1H), 7.46 (m, 2H), 4.44 (t, 1H), 4.17 (d, 2H), 3.28 (m, 2H), 2.80 (m, 2H), 1.68 (d, 2H), 1.58 (m, 1H), 1.10 (m, 2H)	341

Example 23**5-oxo-2-(*t*-Butylcarbonylamino)dibenzothiophene**

5 *m*-Chloroperoxybenzoic acid (50% wt/wt, 770 mg, 2.0 mmol) was added to a solution of 2-(*t*-butylcarbonylamino)dibenzothiophene (Example 2; 566 mg, 2.0 mmol) in 1,2-dichloroethane and the resultant mixture warmed at 80° for 16 hours then cooled and DCM was added (50 ml). The mixture was washed with 2 M sodium hydroxide, water and brine, dried and evaporated *in vacuo* to a brown gum. This was purified by chromatography

10 eluting with 0-2% methanol in DCM. The impure product thus obtained was purified by preparative HPLC (Dynamax C-18 60A column eluted with 40-95% acetonitrile in water plus 0.1% trifluoroacetic acid) to obtain the product (53.5 mg) NMR: 9.59 (s, 1H), 8.35 (d, 1H), 8.03 (d, 1H), 7.97 (d, 1H), 7.93 (d, 1H), 7.80 (dd, 1H), 7.70 (t, 1H), 7.56 (t, 1H), 1.16 (s, 9H); *m/z* 300.

15

Example 24**5,5-Dioxo-2-(4-nitrobenzoxycarbonylamino)dibenzothiophene; and****Example 25****20 5-Oxo-2-(4-nitrobenzoxycarbonylamino)dibenzothiophene**

2-(4-Nitrobenzoxycarbonylamino)dibenzothiophene (Example 1; 364 mg, 1.0 mmol) was suspended in a mixture of glacial acetic acid (5.0 ml) and 30% hydrogen peroxide (565 mg, 5 mmol) and the mixture was warmed at 60°C for 2 hours. The suspension of the title compound was filtered and air dried. TLC: (2:1; EtOAc : isohexane) *R_f* = 0.30.

5-Oxo-2-(4-nitrobenzoxycarbonylamino)dibenzothiophene was also isolated as a by product in this reaction in a 1:5 ratio with the di-oxo compound. TLC: (2:1; EtOAc : isohehexane) Rf = 0.06.


5 Example 26

5,5-Dioxo-2-(isopropylcarbonylamino)dibenzothiophene

5,5-Dioxo-2-aminodibenzothiophene (J Med Chem, (1994), 37 (13), 2085-9; 231 mg, 1.0 mmol) was suspended in a solution of DCM (10 ml) and triethylamine (111 mg, 1.1 mmol). *iso*-Butyrylchloride (106.5 mg, 1.0 mmol) was added dropwise and the resultant mixture was stirred for 16 hours at ambient temperature. The mixture was evaporated *in vacuo* and the residue was purified by chromatography (Bond Elut column) eluting with 0-2.5% methanol in DCM to give the product as a colourless solid (139 mg). NMR: 10.38 (brs, 1H), 8.37 (s, 1H), 7.85-8.00 (m, 3H), 7.80 (d, 1H), 7.74 (d, 1H), 7.63 (t, 1H), 2.66 (m, 1H), 1.13 (d, 6H); m/z 302.

Examples 27-28

Following the procedure of Example 26 using 5,5-dioxo-2-aminodibenzothiophene (J Med Chem, (1994), 37 (13), 2085-9) and the appropriate acid chloride the following compounds were prepared.




Ex	R ¹	NMR	m/z
27		9.67 (s, 1H), 8.39 (s, 1H), 7.98 (d, 1H), 7.94 (d, 1H), 7.86 (m, 2H), 7.78 (t, 1H), 7.64 (t, 1H), 1.24 (s, 9H)	316
28		10.69 (brs, 1H), 8.35 (s, 1H), 7.99 (d, 1H), 7.91 (t, 2H), 7.77 (t, 1H), 7.66 (m, 2H), 4.11 (s, 2H), 3.46 (t, 2H), 2.26 (t, 2H), 1.98 (m, 2H)	357

Example 29**5,5-Dioxo-2-(pyrid-4-ylethylcarbonylamino)dibenzothiophene**

5,5-Dioxo-2-aminodibenzothiophene (J Med Chem, (1994), 37 (13), 2085-9; 115 mg, 0.5 mmol) was dissolved in DMF (2.0 ml) and 3-(pyridin-4-yl)propionic acid (Method 1; 75 mg, 0.5 mmol) was added in one portion followed by EDAC (115 mg, 0.5 mmol) and the resultant mixture was stirred for 72 hours at ambient temperature. It was then poured onto water (50 ml) and extracted with DCM (2 x 50 ml), dried and evaporated *in vacuo*. The residue thus obtained was purified by chromatography (Bond Elut column) eluted with 0-5.0% methanol in DCM to give the product as a colourless solid (31.8 mg). NMR 10.52 (brs, 1H), 8.46 (d, 2H), 8.32 (d, 1H), 7.97 (d, 1H), 7.94 (d, 1H), 7.88 (d, 1H), 7.76 (d, 1H), 7.70 (t, 1H), 7.64 (t, 1H), 7.26 (d, 2H), 2.93 (t, 2H), 2.75 (t, 2H); m/z 365.

Example 30

Following the procedure of Example 29 using 5,5-dioxo-2-aminodibenzothiophene (J Med Chem, (1994), 37 (13), 2085-9) and the appropriate acid the following compound was prepared.



Ex	R ¹	NMR DMSO-d ₆ + d ₄ -acetic acid, 100°C	m/z
30		8.26 (s, 1H), 7.88 (d, 1H), 7.80 (m, 1H), 7.72 (m, 3H), 7.61 (t, 1H), 3.78 (t, 4H), 3.37 (dd, 1H), 3.03 (m, 5H), 1.95 (dd, 1H), 1.26 (d, 3H)	387


Example 31**5,5-Dioxo-2-(1,2,4-triazol-1-ylmethylcarbonylamino)dibenzothiophene**

2-(1,2,4-Triazol-1-ylmethylcarbonylamino)dibenzothiophene (Example 3; 110 mg 0.34 mmol) was suspended in glacial acetic acid (3.0 ml) and 30% hydrogen peroxide was added (0.75 ml). The mixture was warmed to gentle reflux for 30 minutes to give a pale straw-coloured homogeneous solution which was poured onto water (50 ml) and extracted with DCM (2 x 25 ml), dried and evaporated *in vacuo* to leave a pale yellow solid (73 mg).

NMR: 10.94 (brs, 1H), 8.55 (s, 1H), 8.32 (d, 1H), 7.98 (m, 2H), 7.95 (s, 1H), 7.93 (s, 1H), 7.76 (t, 1H), 7.70 (dd, 1H), 7.63 (t, 1H), 5.21 (s, 2H); m/z 341.

Examples 32-37

- 5 Following the procedure of Example 31 using the appropriate dibenzothiophenes the following compounds were prepared.




Ex	R ¹	NMR	m/z	SM
32		10.74 (brs, 1H), 8.35 (d, 1H), 7.9 (m, 3H), 7.8 (t, 1H), 7.7 (dd, 1H), 7.6 (t, 1H), 1.85 (m, 1H), 0.9 (m, 4H)	300	Ex 4
33		10.49 (brs, 1H), 8.35 (d, 1H), 7.95 (d, 1H), 7.9 (d, 1H), 7.85 (d, 1H), 7.8 (t, 1H), 7.7 (dd, 1H), 7.65 (t, 1H), 3.65 (t, 2H), 3.3 (s, 3H), 2.6 (t, 2H)	318	Ex 6
34		10.54 (brs, 1H), 8.35 (s, 1H), 7.9 (m, 3H), 7.8 (t, 1H), 7.7 (m, 2H), 3.4-3.0 (m, 3H), 2.9-2.7 (m, 2H), 2.65 (m, 2H), 2.3 (m, 1H), 1.9 (m, 1H)	392	Ex 7
35		10.64 (brs, 1H), 8.35 (d, 1H), 8.0 (d, 1H), 7.95 (d, 1H), 7.9 (d, 1H), 7.8 (t, 1H), 7.7 (dd, 1H), 7.65 (t, 1H), 4.3 (t, 2H), 4.1 (s, 2H), 3.7 (t, 2H)	359	Ex 8
36		10.43 (brs, 1H), 8.4 (d, 1H), 8.0 (d, 1H), 7.95 (d, 1H), 7.9 (d, 1H), 7.8 (t, 1H), 7.7 (dd, 1H), 7.65 (t, 1H), 3.9 (m, 2H), 3.35 (m, 2H), 2.7 (m, 1H), 1.7 (m, 4H)	344	Ex 9
37		10.64 (brs, 1H), 8.4 (s, 1H), 7.9 (m, 3H), 7.8 (m, 2H), 7.65 (t, 1H), 7.3 (m, 2H), 6.95 (m, 3H), 4.9 (q, 1H), 1.6 (d, 3H)	380	Ex 10

Example 38**5,5-Dioxo-2-[(2-pyrid-4-yl-1-methylethyl)carbonylamino]dibenzothiophene**


2-[(2-Pyrid-4-yl-1-methylethyl)carbonylamino]dibenzothiophene (Example 11; 0.39 g, 1.13 mmol) was dissolved in glacial acetic acid (10 ml) and concentrated sulphuric acid (0.5 ml) at 60°C. 30% aqueous hydrogen peroxide (1.3 ml) was added in one portion and the solution was stirred at 60°C for two hours. The mixture was allowed to cool and then poured into water (50 ml). The mixture was made basic by the addition of solid NaHCO₃ and extracted with DCM (2 x 50 ml). The combined organic layers were washed with brine (50 ml), dried and evaporated *in vacuo* to leave a pale yellow solid. NMR: 8.45 (d, 2H), 8.3 (d, 1H), 8.0 (d, 1H), 7.95 (d, 1H), 7.9 (d, 1H), 7.8 (t, 1H), 7.7 (dd, 1H), 7.65 (t, 1H), 7.25 (d, 2H), 3.0 (m, 2H), 2.7 (m, 1H), 1.15 (d, 3H); m/z 379.

Examples 39-43

Following the procedure of Example 38 using the appropriate dibenzothiophenes the following compounds were prepared.



Ex	R ¹	NMR	m/z	SM
39		8.4 (d, 1H), 8.0 (d, 1H), 7.95 (d, 1H), 7.9 (d, 1H), 7.8 (m, 2H), 7.65 (m, 2H), 7.2 (s, 1H), 6.8 (s, 1H), 4.3 (t, 2H), 2.9 (t, 2H)	354	Ex 14
40		8.5 (s, 1H), 8.45 (d, 1H), 8.0 (d, 1H), 7.9 (m, 2H), 7.85 (d, 1H), 7.8 (m, 2H), 7.65 (t, 1H), 4.5 (t, 2H), 3.0 (t, 2H)	355	Ex 12
41		11.24 (brs, 1H), 8.4 (d, 1H), 8.1 (s, 1H), 8.05 (d, 1H), 8.0 (d, 1H), 7.95 (m, 2H), 7.8-7.6 (m, 5H), 5.5 (s, 2H)	367	Ex 13
42		10.83 (brs, 1H), 8.3 (s, 1H), 8.0 (d, 1H), 7.95 (m, 2H), 7.8 (t, 1H), 7.7 (m, 2H), 4.0 (s, 2H)	297 (M-H) ⁻	Ex 19

Ex	R ¹	m/z	SM
43		316	Ex 17

Example 44

5,5-Dioxo-2-(N'-pyrid-4-ylethyl-N'-methyleureido)dibenzothiophene; and

5 Example 45

5-Oxo-2-(N'-pyrid-4-ylethyl-N'-methyleureido)dibenzothiophene

5,5-Dioxo-2-(4-nitrobenzoxycarbonylamino)dibenzothiophene contaminated with 5-oxo-2-(4-nitrobenzoxycarbonylamino)dibenzothiophene (Examples 24 and 25; 396 mg, 1.0 mmol) was suspended in EtOAc (10 ml) and 4-[2-(methylamino)ethyl]pyridine (272 mg, 2.0 mmol) was added in one portion. The resulting mixture was stirred at ambient temperature for 16 hours then further EtOAc (100 ml) was added and the solution was washed with 1 M sodium hydroxide (2 x 25 ml), water (2 x 25 ml) and brine (25 ml), dried and evaporated *in vacuo*. The resulting mixture was separated by preparative HPLC (Dynamax C-18 60A column eluted with 20-90% acetonitrile in water plus 0.1% trifluoroacetic acid). Two products were obtained:


5-Oxo-2-(N'-pyrid-4-ylethyl-N'-methyleureido)dibenzothiophene (16.5 mg): NMR: 8.71 (s, 1H), 8.67 (d, 2H), 8.17 (s, 1H), 8.02 (d, 1H), 7.85 (m, 2H), 7.72 (d, 2H), 7.67 (t, 1H), 7.56 (m, 2H), 3.69 (t, 2H), 3.05 (t, 2H), 3.01 (s, 3H); m/z 378.

5,5-Dioxo-2-(N'-pyrid-4-ylethyl-N'-methyleureido)dibenzothiophene (82.0 mg): NMR: 8.84 (s, 1H), 8.66 (d, 2H), 8.10 (s, 1H), 7.90 (d, 2H), 7.75 (m, 4H), 7.62 (t, 2H), 3.69 (t, 2H), 3.04 (t, 2H), 3.01 (s, 3H); m/z 394.

Examples 46-49

Following the procedure of Example 44 using 5,5-dioxo-2-(4-nitrobenzoxycarbonyl-amino)dibenzothiophene (Example 24) and the appropriate amine the following compounds were prepared (NB the corresponding sulphoxides were not isolated).

- 64 -



Ex	R ¹	NMR	m/z
46		9.81 (brs, 1H), 8.13 (d, 1H), 7.96 (d, 1H), 7.89 (d, 1H), 7.76 (m, 2H), 7.63 (t, 1H), 7.56 (dd, 1H), 3.42 (t, 2H), 2.96 (s, 3H), 2.46 (t, 2H), (2.24, s, 6H)	360
47		8.80 (s, 1H), 8.20 (s, 1H), 7.92 (m, 2H), 7.77 (m, 2H), 7.64 (m, 2H), 4.02 (m, 1H), 2.05 (s, 3H), 2.02 (d, 2H), 2.15 (s, 3H), 1.95 (t, 2H), 2.14, (m, 2H), 1.51 (d, 2H)	386
48		9.1 (brs, 1H), 8.1 (s, 1H), 7.9 (t, 2H), 7.8 (d, 1H), 7.75 (t, 1H), 7.6 (m, 2H), 3.6 (m, 4H), 3.5 (m, 4H)	345
49		9.1 (s, 1H), 8.2 (s, 1H), 7.9 (t, 2H), 7.8 (m, 2H), 7.6 (m, 2H), 3.5 (m, 4H), 2.3 (m, 4H), 2.2 (s, 3H)	358


Example 50**5 5,5-Dioxo-2-(4,4,4-trifluoro-2-methylbutyrylcarbonylamino)dibenzothiophene**

EDAC (0.307 g, 1.6 mmol) was added to a solution of 2-aminodibenzothiophene (0.299 g, 1.5 mmol), 1-hydroxybenzotriazole (0.216 g, 1.6 mmol) and 4,4,4-trifluoro-2-methylbutanoic acid (0.250 g, 1.6 mmol) in DMF (10 ml). The solution was stirred at room temperature overnight. The solvent was removed *in vacuo* and the residue was shaken in water. The crude product was collected by filtration and dried in air. The crude amide was dissolved in glacial acetic acid (10 ml) at 60°C and hydrogen peroxide (30 wt. % solution in water, 2 ml) was added. The mixture was stirred at 60°C for 2 hours and left to cool to room temperature overnight. The mixture was filtered and the residue was washed with water (10 ml), methanol (5 ml) and finally with ether (3 x 5 ml) to leave a pale yellow






solid (0.147 g). NMR: 10.63 (s, 1H), 8.3 (d, 1H), 8.0 (d, 1H), 7.9 (m, 2H), 7.75 (m, 2H), 7.7 (t, 1H), 2.9 (m, 1H), 2.7 (m, 1H), 2.4 (m, 1H), 1.3 (d, 3H); m/z 370.

Examples 51-65

- 5 Following the procedure of Example 50 using 2-aminodibenzothiophene and the appropriate acid the following compounds were prepared (NB the corresponding sulphides were not isolated).



Ex	R ¹	NMR	MS
51		12.1 (s, 1H), 8.2 (s, 1H), 8.1 (m, 2H), 8.0 (d, 1H), 7.8 (t, 1H), 7.7 (m, 2H), 2.8 (m, 2H), 2.2 (m, 2H), 1.7 (m, 4H), 1.5 (m, 1H)	354 (M-H) ⁻
52		10.68 (s, 1H), 8.4 (s, 1H), 8.0 (d, 1H), 7.9 (m, 2H), 7.7 (m, 3H), 3.3 (m, 1H), 3.0 (dd, 1H), 2.8 (dd, 1H), 1.5 (s, 3H), 1.3 (s, 3H)	370 (M-H) ⁻
53		10.55 (s, 1H), 8.4 (s, 1H), 7.9 (m, 3H), 7.7 (m, 3H), 4.0 (t, 1H), 3.8 (m, 3H), 3.2 (m, 1H), 2.1 (m, 2H)	328 (M-H) ⁻
54		10.42 (s, 1H), 8.35 (s, 1H), 7.9 (m, 3H), 7.8 (t, 1H), 7.7 (dd, 1H), 7.65 (t, 1H), 3.6 (m, 2H), 2.8 (m, 1H), 1.7-1.4 (m, 4H), 1.2 (s, 3H), 1.15 (s, 3H)	372
55 ¹		10.81 (s, 1H), 8.3 (s, 1H), 8.0 (d, 1H), 7.9 (m, 2H), 7.8 (t, 1H), 7.65 (m, 2H), 4.3 (s, 2H), 2.75 (s, 4H)	371
56		10.59 (s, 1H), 8.3 (s, 1H), 7.9 (m, 3H), 7.8 (t, 1H), 7.7 (m, 2H), 7.3 (m, 2H), 7.2 (m, 3H), 4.4 (s, 2H), 3.5 (m, 1H), 3.2 (m, 2H), 2.65 (m, 2H)	433

57 ²		10.7 (s, 1H), 8.35 (s, 1H), 8.0 (d, 1H), 7.9 (m, 2H), 7.8 (t, 1H), 7.7 (m, 2H), 7.4 (m, 5H), 5.7 (d, 1H), 3.1 (dd, 1H), 2.9 (dd, 1H)	418 (M-H) ⁻
58		10.54 (s, 1H), 8.3 (s, 1H), 7.9 (m, 3H), 7.8 (t, 1H), 7.7 (m, 2H), 3.7 (t, 2H), 2.6 (m, 6H)	385
59 ³		10.15 (s, 1H), 8.4 (d, 1H), 7.9 (m, 2H), 7.85 (d, 1H), 7.75 (t, 1H), 7.65 (m, 2H), 4.8 (d, 1H), 4.0 (m, 1H), 2.55 (d, 1H), 2.4 (bs, 1H), 2.0 (m, 2H), 1.4 (m, 2H), 1.1 (m, 3H)	370
60 ⁴		10.64 (s, 1H), 8.3 (d, 1H), 8.0 (d, 1H), 7.9 (m, 2H), 7.8 (t, 1H), 7.7 (m, 2H), 7.25 (m, 5H), 4.2 (d, 1H), 4.1 (d, 1H), 3.3 (m, 2H), 3.1 (dd, 1H), 2.7 (m, 1H), 2.6 (m, 1H), 2.0 (m, 1H), 1.7 (m, 1H)	447
61 ⁵		NMR (400 MHz) 12.2 (s, 1H), 10.7 (s, 1H), 8.35 (d, 1H), 8.05 (d, 1H), 7.95 (m, 2H), 7.8 (td, 1H), 7.75 (dd, 1H), 7.7 (td, 1H), 7.35 (d, 1H), 6.95 (d, 1H), 4.9 (s, 2H)	384



¹ Acid: Al-Azhar J. Pharm. Sci., 1998, 21, 133-141



² Acid: Arch. Immunol. Ther. Exp., 1968, 16 (1), 155-172

³ Acid: J. Am. Chem. Soc., 1974, 96 (20), 6492-6498

⁴ Acid: Method 4

5 ⁵ Acid: Arm. Khim. Zh., 1971, 24 (4), 350-353

Ex	R ¹	MS	Acid Preparation Reference
62		367	Rev. Roum. Chim., 1988, 33 (7), 729-739
63		369	Method 6

64		357	J. Org. Chem., 1980, 45(5), 810-814
65		433	J. Org. Chem., 1969, 34(10), 3187-3189

Example 665,5-Dioxo-2-(3-methyl-3-pyridin-4-ylpropionamido)dibenzothiophene

EDAC (0.500 g, 2.6 mmol) was added to a solution of 2-aminodibenzothiophene (0.470 g, 2.4 mmol), 1-hydroxybenzotriazole (0.351 g, 2.6 mmol) and 3-methyl-3-pyridin-4-ylpropanoic acid (Method 7; 0.429 g, 2.6 mmol) in DMF (10 ml). The solution was then stirred at room temperature overnight. The solvent was removed *in vacuo* and the residue was then shaken in water (50 ml) and DCM (50 ml). The organic layer was separated, washed with brine (50 ml), dried and then evaporated *in vacuo* to leave a light brown solid. The crude amide was dissolved in a mixture of glacial acetic acid (15 ml) and concentrated sulphuric acid (0.5 ml) at 60°C and hydrogen peroxide (30 wt. % solution in water, 3 ml) was added. The mixture was stirred at 60°C for 2 hours and left to cool to room temperature overnight. The mixture was poured into water (50 ml) and made basic by the addition of potassium carbonate. The mixture was extracted into DCM (50 ml), the organic layer was separated and washed with brine (50 ml), dried and evaporated *in vacuo* to leave a light brown solid. The crude product was triturated with methanol (10 ml) to leave a pale yellow solid (0.496 g). NMR: 10.47 (s, 1H), 8.3 (d, 1H), 7.9 (m, 3H), 7.75 (t, 1H), 7.65 (m, 2H), 3.3 (m, 1H), 2.7 (m, 2H), 1.3 (d, 3H); m/z 379.

Example 671-Cyano-5,5-dioxo-2-pivaloylaminodibenzothiophene

A mixture of 1-cyano-2-pivaloylaminodibenzothiophene (Example 68; 0.034 g, 0.11 mmol) glacial acetic acid (2 ml) and 30% hydrogen peroxide (0.2 ml) was heated at 80°C for 4 hours. The mixture was cooled and poured into water. The precipitate was filtered and

washed extensively with water to give the title compound. NMR 9.95 (s, 1H), 8.57 (d, 1H), 8.32 (d, 1H), 8.11 (d, 1H), 7.95 (t, 1H), 7.82 (t, 1H), 7.75 (d, 1H), 1.28 (s, 9H); m/z 341.

Example 68

5 1-Cyano-2-pivaloylaminodibenzothiophene

A mixture of 2-amino-1-cyanodibenzothiophene (J. Heterocycl. Chem., 1977, 14(7), 1209-1214; 0.043 g, 0.19 mmol), triethylamine (0.04 g, 0.4 mmol) and pivaloyl chloride (0.05 g, 0.4 mmol) in DCM (2 ml) was stirred at ambient temperature for 24 hours. The resultant mixture was extracted with water and the organic phase reduced to a residue which was
10 triturated with ether/iso-hexane to give the title compound. NMR: 9.78 (s, 1H), 8.81 (m, 1H), 8.38 (d, 1H), 8.16 (m, 1H), 7.58 (m, 2H), 7.48 (d, 1H), 1.18 (s, 9H); m/z 307 (M-H)⁺.

Example 69

3-Methyl-5,5-dioxo-2-pivaloylaminodibenzothiophene

15 3-Bromo-5,5-dioxo-2-pivaloylaminodibenzothiophene (Example 70; 0.1 g, 0.254 mmol) was suspended in dry THF (2 ml) under argon, cooled to -70°C and 1.7 M *tert*-butyllithium in pentane (0.49 ml, 0.838 mmol) was added dropwise over 5 minutes. The mixture was allowed to warm to -30°C and the suspension slowly dissolved as the solution darkened. The mixture was then re-cooled to -70°C and iodomethane (0.17 ml, 0.279 mmol)
20 added and stirred for 30 minutes. Saturated ammonium acetate (5 ml) and DCM (25 ml) were added and the organic phase washed with water and dried. The mixture was evaporated *in vacuo* and the residue purified by preparative HPLC (Dynamax C-18 60A column eluted with 30-90% acetonitrile in water plus 0.1% trifluoroacetic acid) to give the product as a colourless solid. NMR: 9.16 (s, 1H), 8.13 (d, 1H), 7.99 (s, 1H), 7.92 (d, 1H), 7.86 (s, 1H), 7.76 (t, 1H),
25 7.61 (t, 1H), 2.28 (s, 3H), 1.26 (s, 9H); m/z 330.

Example 70

3-Bromo-5,5-dioxo-2-pivaloylaminodibenzothiophene

2-Amino-3-bromo-5,5-dioxidodibenzothiophene (DE 2638081; 0.309 g, 1.0 mmol)
30 was suspended in a solution of DCM (5 ml) and triethylamine (0.111 g, 1.1 mmol). Pivaloyl chloride (0.132 g, 1.1 mmol) was added dropwise and the resultant mixture stirred for 5 days at ambient temperature. The mixture was reduced to dryness and the residue was triturated

with methanol. NMR: 9.25 (s, 1H), 8.42 (s, 1H), 8.31 (s, 1H), 8.11 (d, 1H), 7.98 (d, 1H), 7.80 (t, 1H), 7.68 (t, 1H), 1.28 (s, 9H); m/z 394/396.

Example 71

5 1-Methyl-5,5-dioxo-2-(tetrahydrofur-3-ylcarbonylamino)dibenzothiophene

1-Methyl-2-(tetrahydrofur-3-ylcarbonylamino)dibenzothiophene (Example 72; 436 mg, 1.4 mmol) was stirred in glacial acetic acid (10 ml) and hydrogen peroxide (30% w/v, 2 ml) at 120°C for 30 mins. The reaction mixture was cooled and the precipitate was filtered and washed with water to leave the product as a yellow crystalline solid. NMR: 9.96 (brs, 1H), 8.23 (d, 1H), 8.00 (d, 1H), 7.80 (m, 2H), 7.64 (m, 2H), 3.96 (m, 1H), 3.76 (m, 3H), 3.27 (m, 1H), 2.55 (s, 3H), 2.15 (q, 2H); m/z 344.

Example 72

1-Methyl-2-(tetrahydrofur-3-ylcarbonylamino)dibenzothiophene

15 2-Amino-1-methyldibenzothiophene (Method 10; 0.337 g, 1.58 mmol), 4-(dimethylamino)pyridine (0.213 g, 1.74 mmol), EDAC (0.334 g, 1.74 mmol) and tetrahydro-3-furoic acid (0.167 ml, 1.74 mmol) were stirred in DCM (20 ml) at reflux under an argon atmosphere for 64 h. The product was purified by flash chromatography eluting with DCM - 5% methanol/DCM. Evaporation of the appropriate fractions gave the product as a
20 white solid. NMR: 9.77 (brs, 1H), 8.44 (m, 1H), 8.04 (m, 1H), 7.82 (d, 1H), 7.52 (m, 2H), 7.36 (d, 1H), 4.00 (m, 1H), 3.76 (m, 3H), 3.27 (m, 1H), 2.68 (s, 3H), 2.15 (q, 2H); m/z 313.

Example 73

5,5-Dioxo-2-(6-methylpyridazin-3-ylamino)dibenzothiophene

25 Hydrogen peroxide (100 vols, 1 ml) was added to 2-(6-methylpyridazin-3-ylamino)dibenzothiophene (Example 74; 300 mgs, 1.03 mmol) in acetic acid (3 ml) and conc. sulphuric acid (0.1 ml) and the mixture was heated to 60°C for 40 minutes. On cooling to room temperature the mixture was diluted with DCM: MeOH (1:19) washed with aqueous potassium carbonate, dried over sodium sulphate and concentrated.
30 Chromatography (eluent gradient of DCM to DCM: MeOH(1:1)) gave the product as an off white solid (0.131 g). NMR: 9.87 (brs, 1H), 8.48 (brs, 1H), 8.00-7.70 (m, 5H), 7.65 (t, 1H), 7.45 (d, 1H), 7.20 (d, 1H), 2.52 (s, 3H); m/z 324.28.

Example 73**2-(6-Methylpyridazin-3-ylamino)dibenzothiophene**

- A mixture of 2-aminodibenzothiophene (0.90 g, 4.52 mM),
3-chloro-6-methyl-pyridazine (0.58 g, 4.52 mM), sodium t-butoxide (0.61 g, 6.33 mM),
5 tris(dibenzylidene-acetone)dipalladium(0) (0.083 g, 0.09 mM) and
S-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (0.113 g, 0.18 mM) in toluene (23 ml) was
heated to 80°C for 18 hours under an inert atmosphere. On cooling to room temperature the
mixture was diluted with DCM, washed with aqueous potassium carbonate, dried over sodium
sulphate and concentrated. Chromatography (eluent gradient of DCM to EtOAc then MeOH:
10 EtOAc (1:9)) gave the product as an off white solid (0.456 g). NMR: 9.38 (brs, 1H), 8.81 (brs,
1H), 8.18 (m, 1H), 8.01 (m, 1H), 7.94 (d, 1H), 7.75 (dd, 1H), 7.50 (m, 1H), 7.36 (d, 1H), 7.14
(d, 1H), 2.50 (s, 3H); m/z 292.29.


Example 74









- 15 **5,5-Dioxo-2-(N'-pyrid-4-ylmethyl-N'-ethylureido)dibenzothiophene**










- 5,5-Dioxo-2-(4-nitrobenzoxycarbonylamino)dibenzothiophene (Example 24; 0.396 g,
1 mmol) was dissolved in EtOAc (10 ml) and treated with 4-(ethylaminomethyl)pyridine
(0.150 g, 1.1 mmol), triethylamine (0.202 g, 2 mmol) and 4-(dimethylamino)pyridine (0.006
mg, 0.05 mmol). The reaction was stirred for 18 hours and the suspended solids collected,
20 washed with 1 M aq. NaOH solution, water and then ether to leave the product as a colourless
solid. NMR: 9.00 (s, 1H), 8.54 (m, 2H), 8.27 (s, 1H), 7.93 (m, 2H), 7.81 (m, 1H), 7.75 (m,
2H), 7.65 (m, 1H), 7.28 (m, 2H), 4.63 (s, 2H), 3.43 (q, 2H), 1.12 (t, 3H); m/z 394.42.

Examples 75-192









- 25 Following the procedure of Example 74 using
5,5-Dioxo-2-(4-nitrobenzoxycarbonylamino)dibenzothiophene (Example 24) and the
appropriate amine the following compounds were prepared.





















Ex	R ¹	N. M. R.	MS
75		9.04 (s, 1H), 8.21 (d, 1H), 7.93 (t, 2H), 7.79 (d, 1H), 7.75 (d, 1H), 7.64 (m, 2H), 4.15 (d, 2H), 3.26 (m, 2H), 2.82 (t, 2H), 1.70 (d, 2H), 1.60 (m, 1H), 1.08 (m, 2H)	373
76		9.03 (s, 1H), 8.54 (d, 1H), 8.48 (dd, 1H), 8.25 (s, 1H), 7.96 (d, 1H), 7.92 (d, 1H), 7.83 (d, 1H), 7.72 (m, 3H), 7.62 (t, 1H), 7.37 (dd, 1H), 4.61 (s, 2H), 3.02 (s, 3H)	380
77		9.12 (s, 1H), 8.48 (d, 2H), 8.15 (s, 1H), 7.97 (d, 1H), 7.90 (d, 1H), 7.7.9 (m, 2H), 7.66 (m, 1H), 7.52 (m, 1H), 7.28 (d, 2H), 6.48 (t, 1H), 3.44 (m, 2H), 2.81 (t, 2H)	380
78		9.13 (s, 1H), 8.16 (s, 1H), 7.99 (d, 1H), 7.90 (d, 1H), 7.77 (m, 2H), 7.53 (m, 2H), 7.53 (d, 1H), 7.19 (s, 1H), 6.90 (s, 1H), 6.52 (t, 1H), 4.00 (t, 2H), 3.10 (m, 2H), 1.89 (t, 2H)	383
79		8.10 (s, 1H), 7.90 (m, 3H), 7.81 (m, 1H), 7.78 (m, 1H), 7.74 (m, 1H), 7.46 (d, 1H), 4.0 (m, 1H), 3.57 (m, 1H), 3.29 (m, 1H), 2.75 (m, 3H), 2.60 (m, 2H), 2.50 (m, 1H), 2.02 (m, 1H), 1.65 (m, 7H)	412
80		9.10 (s, 1H), 8.20 (s, 1H), 7.93 (m, 2H), 7.80 (m, 2H), 7.64 (m, 2H), 7.33 (s, 1H), 6.88 (s, 1H), 4.10 (m, 1H), 4.05 (m, 1H), 2.93 (m, 1H), 2.84 (m, 1H), 2.30 (m, 1H), 1.92 (m, 1H), 1.68 (m, 1H), 1.52 (m, 1H), 1.42 (m, 1H)	386
81	 trans	8.93 (s, 1H), 8.15 (s, 1H), 7.98 (d, 1H), 7.90 (d, 1H), 7.75 (m, 1H), 7.64 (m, 1H), 6.31 (d, 1H), 4.50 (d, 1H), 3.42 (m, 1H), 1.83 (m, 4H), 1.23 (m, 4H)	373
82		9.0 (s, 1H), 8.2 (d, 1H), 7.9 (t, 2H), 7.8 (m, 2H), 7.6 (m, 2H), 4.3 (t, 1H), 4.1 (m, 1H), 3.45 (q, 2H), 1.7 (m, 3H), 1.4 (q, 2H), 1.1 (m, 2H)	387









83		8.79 (s, 1H), 8.19 (s, 1H), 7.93 (m, 2H), 7.78 (m, 2H), 7.65 (m, 2H), 7.27 (m, 4H), 7.18 (m, 1H), 3.79 (t, 2H), 2.98 (s, 3H), 2.84 (t, 2H)	393
84		9.10 (s, 1H), 8.11 (d, 1H), 7.92 (m, 2H), 7.78 (m, 2H), 7.65 (m, 2H), 4.07 (m, 2H), 2.98 (m, 2H), 2.42 (m, 1H), 1.89 (m, 2H), 1.50 (m, 2H)	387
85		9.18 (s, 1H), 8.14 (d, 1H), 7.96 (m, 1H), 7.90 (m, 1H), 7.76 (m, 2H), 7.64 (t, 1H), 7.52 (dd, 1H), 6.35 (t, 1H), 3.57 (t, 2H), 3.24 (m, 2H), 2.40 (m, 6H)	388
86 ¹		9.20 (s, 1H), 8.20 (d, 1H), 7.92 (m, 2H), 7.79 (m, 2H), 7.65 (m, 2H), 3.53 (m, 4H), 3.47 (t, 2H), 3.00 (s, 3H), 2.44 (m, 4H)	402
87 ²		8.62 (s, 1H), 8.13 (s, 1H), 7.93 (m, 2H), 7.77 (m, 2H), 7.62 (m, 2H), 4.52 (d, 1H), 3.82 (m, 1H), 3.39 (m, 2H), 2.00 (m, 2H), 1.84 (m, 2H), 1.55 (m, 2H), 1.33 (m, 1H), 1.28 (d, 6H)	415
88 ³		8.75 (s, 1H), 8.47 (d, 2H), 8.24 (s, 1H), 7.93 (m, 2H), 7.80 (m, 1H), 7.70 (m, 2H), 7.66 (m, 1H), 7.28 (d, 2H), 3.62 (t, 2H), 2.90 (t, 2H), 2.65 (m, 1H), 0.91 (m, 2H), 0.72 (m, 2H)	420
89		8.79 (s, 1H), 8.51 (d, 2H), 8.27 (s, 1H), 7.92 (d, 2H), 7.78 (m, 3H), 7.63 (t, 1H), 7.35 (d, 2H), 3.94 (t, 1H), 3.67 (t, 1H), 3.44 (m, 3H), 2.32 (m, 1H), 2.01 (m, 1H)	406
90		8.79 (s, 1H), 8.57 (s, 1H), 8.44 (d, 1H), 8.28 (s, 1H), 7.91 (d, 2H), 7.78 (m, 4H), 7.61 (t, 1H), 7.35 (m, 1H), 3.96 (t, 1H), 3.69 (t, 1H), 3.46 (m, 3H), 2.32 (m, 1H), 2.02 (m, 1H)	406
91 ⁴		9.3 (s, 1H), 8.2 (d, 1H), 7.95 (m, 2H), 7.85 (d, 1H), 7.75 (t, 1H), 7.65 (m, 2H), 3.65 (m, 4H), 2.0 (m, 4H)	379










¹ Amine: Tetrahedron, 1992, 48(11), 1999-2012² Amine: Tetrahedron Lett., 1995, 36(10), 1709-1712³ Amine: Method 12⁴ Amine: Chem. Pharm. Bull., 1993, 41(11), 1971-1986











Ex	R ¹	MS	Amine Preparation Reference
92		331	WO 0109138
93		341	Commercially Available
94		409	Method 13
95	 trans	403	Method 17
96		425	US 4798892
97		419	Commercially Available
98		347	J. Org. Chem., 1990, 55(24), 5935-5936.
99		415	WO 9418182










100		451	Heterocycles, 2000, 53(4), 797-804.
101		394	Commercially Available
102		464	Commercially Available
103		428	DE 3440195
104		416	Commercially Available
105		420	Bull. Soc. Chim. Fr., 1996, 133(4), 369-379.
106		442	Commercially Available
107		421	Commercially Available
108		377	Commercially Available










109		347	Commercially Available
110		369	Commercially Available
111		381	Commercially Available
112		373	Commercially Available
113		387	Commercially Available
114		415	Collect. Czech. Chem. Commun., 1986, 51(11), 2598-2616
115		416	WO 9727188
116		429	JP 03188030
117		429	Commercially Available










118		401	Commercially Available
119		393	Commercially Available
120		361	Commercially Available
121		421	Commercially Available
122		432	Indian J. Chem., 1972, 10(7), 766-777.
123		436	DE 3905364
124		493	Method 19
125		422	WO 9811128









126		382	Method 21
127		416	Commercially Available
128		442	WO 0039114
129		424	DE 2362568
130		373	J. Org. Chem., 1987, 52(1), 15-18.
131		455	Commercially Available
132		386	Commercially Available
133		388	Tetrahedron Lett., 1995, 36(8), 1267-1270.
134	 (R)-enantiomer	372	J. Med. Chem., 1985, 28(11), 1558-1564.









135	 <chem>CC(=O)N(C)CC(O)c1ccc(O)cc1</chem>	425	Commercially Available
136	 <chem>CC(=O)N1CCCC1C(=O)OCc2ccccc2</chem>	463	Commercially Available
137	 <chem>CC(=O)N(C)C1CCNC1</chem>	372	Commercially Available
138	 <chem>CC(=O)N1CCNC1Cc2ccccc2</chem>	434	Commercially Available
139	 <chem>CC(=O)N1CCCC1(C)(C)c2ccccc2</chem>	433	J. Med. Chem. (1998), 41(26), 5320-5333
140	 <chem>CC(=O)NCCCCN(C)Cc1ccccc1</chem>	422	Commercially Available
141	 <chem>CC(=O)NCCCN#Cc1ccsc1</chem>	424	Commercially Available
142	 <chem>CC(=O)NCCCN#Cc1ccccc1</chem>	418	Commercially Available
143	 <chem>CC(=O)N(C)Cc1cc(C)nc(C)c1</chem>	394	Acta Chim. Hung. (1989), 126(4), 441-54.
144	 <chem>CC(=O)N1CCCC1(C)(C)Cc2ccccc2</chem>	462	Commercially Available





145	 (R)-enantiomer	448	Commercially Available
146		436	DE 2625468
147		478	Commercially Available
148		424	WO 9109857
149		438	J. Med. Chem., 1972, 15(12), 1321-1324.
150		437	EP 421762
151		400	US 5914405
152		384	Method 23
153		375	Chem. Phys. Lipids, 1992, 61(2), 199-208.

154		456	WO 9806705
155		414	Commercially Available
156		341	Commercially Available
157		357	Commercially Available
158		357	Commercially Available
159		397	Commercially Available
160		429	Commercially Available
161		401	Khim. -Farm. Zh., 1984, 18(12), 1445-1448.
162		385	Commercially Available

163		355	Heterocycles, 1982, 19(11), 2155-2182.
164		369	Commercially Available
165		355	Commercially Available
166		383	Commercially Available
167		419	J. Med. Chem., 1991, 34(11), 3212-3228.
168		407	Commercially Available
169		407	Commercially Available
170		468	Commercially Available
171		430	Commercially Available

172		475	Commercially Available
173		450	Commercially Available
174		357	Commercially Available
175		379	Commercially Available
176		413	Commercially Available
177		413	Commercially Available
178		452 (M-H) ⁺	Commercially Available
179		448 (M-H) ⁺	Commercially Available

180	 <chem>COc1ccc(cc1)N2CCN(CC2)C(=O)N</chem>	448 (M-H) ⁻	Commercially Available
181	 <chem>Clc1ccc(cc1)N2CCN(CC2)C(=O)N</chem>	452 (M-H) ⁻	Commercially Available
182	 <chem>O=C1NC(=O)N2C(=N1)C3=CC=CC=C3C2N4CCN(CC4)C(=O)N</chem>	473	Commercially Available
183	 <chem>Clc1cccc(c1)CN2CCN(CC2)C(=O)N</chem>	413	Commercially Available
184	 <chem>C1CCN(CC1)C(=O)NCC2CCCCC2</chem>	385	Commercially Available
185	 <chem>COc1cc(OC)ccc1CN2CCN(CC2)C(=O)N</chem>	453	Commercially Available
186	 <chem>C=C1CC(S(=O)(=O)C1)N2CCN(CC2)C(=O)N</chem>	433	Commercially Available
187	 <chem>c1ccccc1N2CCN(CC2)C(=O)N</chem>	419	Commercially Available

188		445	Commercially Available
189		397	J. Chem. Soc., Perkin Trans. 1, 1998, 16, 2527-2532.
190		437	Commercially Available
191		475	Khim. -Farm. Zh., 1987, 21(7), 808-811.

Example 1925,5-Dioxo-2-[4-(morpholinocarbonyl)piperidinocarbonylamino]dibenzothiophene

5,5-Dioxo-2-(4-carboxypiperidinocarbonylamino)dibenzothiophene (Example 84; 75 mg, 0.19 mmol), DMF (2.5 ml), morpholine (16.5 mg, 0.19 mmol), EDAC (44 mg, 0.23 mmol), 1-hydroxybenzotriazole hydrate (25.65 mg, 0.19 mmol) were stirred together for 18 hours. The solvent was removed under vacuum and the residue partitioned between H₂O / EtOAc. The organic layers were washed with water, saturated aqueous NaHCO₃, then brine, and then dried, filtered and evaporated to an amorphous solid. NMR: 9.08 (s, 1H) 8.21 (d, 1H), 7.93 (m, 2H), 7.79 (m, 2H), 7.65 (m, 2H), 4.14 (m, 2H), 3.52 (m, 6H), 3.45 (m, 2H), 2.92 (m, 3H), 1.68 (m, 2H), 1.53 (m, 2H); m/z 456.73.

Example 1935,5-Dioxo-2-[4-(N,N-dimethylcarbamoyl)piperidinocarbonylamino]dibenzothiophene

The title compound was prepared by the procedure of Example 192 using dimethylamine (1 equivalent) in place of morpholine. NMR: 9.08 (s, 1H), 9.19 (d, 1H), 7.93 (m, 2H), 7.77 (m, 2H), 7.65 (m, 2H), 4.15 (m, 2H), 3.07 (s, 3H), 2.94 (m, 3H), 2.83 (s, 3H), 1.68 (m, 2H), 1.52 (m, 2H); m/z 414.71.

Reference Examples

The following compounds are provided as a further feature of the invention.

Reference Example 1

- 5 5,5-Dioxo-2-[(2,2,2-trifluoro-1-hydroxy-1-methylethyl)carbonylamino]dibenzothiophene

J Med Chem, (1997), 40 (6), 1048.

Reference Example 2**2-Acetamidodibenzothiophene**

- 10 The title compound was prepared following the procedure of Example 2 using 2-aminodibenzothiophene (Bull. Soc. Chim. Fr. (1996), 133(6), 597-610) and acetyl chloride. NMR: 10.13 (brs, 1H), 8.6 (d, 1H), 8.15 (m, 1H), 8.0 (m, 1H), 7.9 (d, 1H), 7.6 (dd, 1H), 7.5 (m, 2H), 2.1 (s, 3H); m/z 242.

- 15 **Reference Example 3**

5,5-Dioxo-2-acetamidodibenzothiophene

The title compound was prepared following the procedure of Example 31 using 2-acetamidodibenzothiophene (Reference Example 2) as the starting material. NMR: 10.48 (brs, 1H), 8.3 (d, 1H), 7.9 (m, 3H), 7.8 (t, 1H), 7.7 (dd, 1H), 7.6 (t, 1H), 2.1 (s, 3H); m/z 274.

20

Preparation of Starting Materials

The starting materials for the Examples above are either commercially available or are readily prepared by standard methods from known materials. For example the following reactions are illustrations but not limitations of the preparation of some of the starting

- 25 materials used in the above reactions.

Method 1**3-(Pyridin-4-yl)propanoic acid**

- 30 To a solution of ethyl 3-pyridin-4-ylpropanoate (Method 2; 103.1 g, 576 mmol) in water (400 ml) and ethanol (20 ml) at room temperature was added potassium hydroxide (60 g, 1600 mmol). After 18 hours hydrochloric acid (100 ml) was added to give a white solid. Yield 62.8 g (73%). NMR 8.38 (d, 2H), 7.21 (d, 2H), 2.70 (t, 2H), 2.52 (t, 2H); m/z 152.2.

Method 2**Ethyl 3-pyridin-4-ylpropanoate**

Ethyl (E)-3-pyridin-4-ylprop-2-enoate (Method 3; 102.3 g, 576 mmol) in methanol (300 ml) was hydrogenated using palladium on carbon 5% (9.0 g) under atmospheric pressure 5 hydrogen for 72 hours. The catalyst was filtered off through diatomaceous earth and the filtrate concentrated to give a yellow oil. Yield 103.1 g (99%). NMR (CDCl₃) 8.50 (d, 2H), 7.15 (d, 2H), 4.12 (q, 2H), 2.95 (t, 2H), 2.64 (t, 2H), 1.21 (t, 3H); m/z 180.4.

Method 310 **Ethyl (E)-3-pyridin-4-ylprop-2-enoate**

To a solution of 4-pyridinecarboxaldehyde (67 ml, 700 mmol) and triethyl phosphonacetate (152 ml, 770 mmol) in THF (200 ml) at room temperature was added lithium hydroxide (32.4 g, 770 mmol). After 18 hours ether (500 ml) was added and the solution was washed with sodium hydrogen carbonate, brine and concentrated to give a white solid. Yield 15 102.1 g (83%). NMR: 8.62 (d, 2H), 7.60 (d, 1H), 7.35 (d, 2H), 6.59 (d, 1H), 4.30 (q, 2H), 1.35 (t, 3H); m/z 178.3.

Method 4**1-Carboxymethyl-3-benzyl-2-pyrrolidinone**

20 A mixture of 3-benzyl-1-(ethoxycarbonylmethyl)-2-pyrrolidinone (Method 5; 0.70 g, 2.68 mmol) and 2 M NaOH (4 ml) was stirred in ethanol (10 ml) for 24 hours. The mixture was evaporated *in vacuo* and the residue acidified. The product was collected by filtration and dried under high vacuum. NMR (90 MHz, CDCl₃) 7.9 (s, 1H), 7.2 (m, 5H), 4.1 (d, 1H), 3.3 (m, 2H), 2.25 (m, 2H), 2.3-1.6 (m, 4H).

25

Method 5**3-Benzyl-1-(ethoxycarbonylmethyl)-2-pyrrolidinone**

NaH (50%, 0.214 g, 4.5 mmol) was added to a solution of 3-benzyl-2-pyrrolidinone (Synthesis, 1996, 8, 941-948; 0.68 g, 3.9 mmol) in THF (15 ml). The solution was stirred at 30 room temperature for 30 minutes. Ethyl bromoacetate (0.48 ml, 4.3 mmol) was added and the mixture stirred for 12 hours. The solvent was removed *in vacuo* and the residue purified by chromatography eluting with 50% EtOAc in hexane to give the product as a light yellow oil.

- 87 -

NMR (90 MHz, CDCl₃) 7.2 (m, 5H), 4.2 (m, 3H), 3.25 (m, 1H), 2.75 (m, 2H), 2.3-1.6 (m, 4H), 1.3 (t, 3H).

Method 6

5 1-(2-Carboxypropyl)-1,2,4-triazole

A mixture of 1,2,4-triazole (27.6 g) and methacrylic acid (34.4 g) in pyridine (40 ml) was heated at 140°C for 6 hours. The precipitate was dissolved in hot hydrochloric acid (1%, 400 ml). Charcoal was added and the mixture was filtered hot. The solution was concentrated *in vacuo* and the colourless solid collected by filtration and washed with cold water. NMR (90
10 MHz): 8.48 (s, 1H), 7.98 (s, 1H), 4.32 (m, 2H), 2.94 (m, 1H), 1.01 (d, 3H).

Method 7

3-Methyl-3-pyridin-4-ylpropanoic acid

Sodium hydroxide (0.19 g, 4.8 mmol) was added to a solution of ethyl
15 3-methyl-3-pyridin-4-ylpropanoate (Method 8; 0.501 g, 2.6 mmol) in MeOH (10 ml) and water (5 ml). The mixture was stirred at room temperature for 18 hours. The solvent was removed *in vacuo* and the residue shaken with 1 M hydrochloric acid (4.8 ml). The solvent was removed *in vacuo* to leave a colourless solid. M/z 164 (M-H).

20 Method 8

Ethyl 3-methyl-3-pyridin-4-ylpropanoate

To a solution of ethyl 3-methyl-3-pyridin-4-ylprop-2-enoate (Method 9; 8.4 g, 44 mmol) in ethanol (150 ml) was added 5% palladium on charcoal. The mixture was stirred under an atmosphere of hydrogen for 18 hours. The mixture was filtered through
25 diatomaceous earth and the filtrate evaporated to leave a colourless oil. The crude product was purified by chromatography eluting with 30-50% EtOAc in hexane to give the product as a colourless oil. M/z 194.

Method 9

30 Ethyl 3-methyl-3-pyridin-4-ylprop-2-enoate

4-Acetyl pyridine (9 ml, 83 mmol) was added to a cooled solution of triethylphosphonoacetate (16.5 ml, 83 mmol) and lithium bis(trimethylsilyl)amide (1 M in

- 88 -

THF, 90 ml, 90 mmol) in THF (100 ml) at 0°C. The mixture was allowed to warm to room temperature and poured into water. The mixture was extracted with EtOAc. The organic layers were dried and evaporated *in vacuo*. The crude product was purified by chromatography eluting with 30-50% EtOAc in hexane to give the product as a yellow oil.

5 M/z 192.

Method 10

2-Amino-1-methyldibenzothiophene

1-Methyl-2-nitrodibenzothiophene (Method 11; 0.855 g, 3.52 mmol) was stirred in
10 ethanol (50 ml) and EtOAc (50 ml) under an argon atmosphere. 10% Palladium on carbon (0.09 g) was added to the solution and the reaction mixture was stirred under a hydrogen atmosphere at room temperature for 3 h. The mixture was filtered through diatomaceous earth and washed with EtOAc before the filtrate was evaporated *in vacuo*. The crude product was purified by flash chromatography eluting with 50% DCM/iso-hexane - DCM. Evaporation of
15 the appropriate fractions left the product as a colourless solid. NMR: 8.39 (m, 1H), 7.92 (m, 1H), 7.50 (m, 1H), 7.42 (m, 2H), 6.92 (d, 1H), 4.98 (brs, 2H), 2.60 (s, 3H); m/z 215.

Method 11

1-Methyl-2-nitrodibenzothiophene

20 To a stirred suspension of 2-nitrodibenzothiophene (2.20 g, 9.61 mmol) in dry THF (100 ml) at -15°C was slowly added methyl magnesium chloride (3 M solution in THF; 9.6 ml, 29 mmol). After stirring at this temperature for 5.5 h,
2,3-dichloro-5,6-dicyano-1,4-benzoquinone (6.98 g, 30.75 mmol) was added portion wise, keeping the temperature below -10°C. The mixture was allowed to warm to room temperature
25 and stirred for 16h. The reaction mixture was diluted with water and extracted into DCM. The organic layers were further washed with water, dried and purified by chromatography (eluent - 50% DCM /isohexane) to give the title compound as a yellow solid. NMR: 8.52 (m, 1H), 8.12 (m, 2H), 7.94 (d, 1H), 7.61 (m, 2H), 2.92 (s, 3H).

Method 12

4-(2-cyclopropylaminoethyl)pyridine

4-Vinyl pyridine (50 mmol), glacial acetic acid (3 ml), cyclopropylamine (4.28 g, 75 mmol) and water (12.8 ml) were heated at 50°C for 22 hours. The crude reaction products

were added to a vigorously stirred mixture of ether and saturated aqueous NaHCO_3 . After partitioning, the organic phase was washed with water, brine and dried. Filtered and solvents removed under vacuum. The residue was further distilled under vacuum (Kugelrohr) and the product amine collected. NMR (CDCl_3) 8.45 (m, 2H), 7.05 (m, 2H), 2.93 (m, 2H), 2.73 (m, 2H), 2.07 (m, 1H), 1.37 (m, 2H), 1.25 (m, 2H); m/z 163.17.

Method 13

***N*-Methyl-*N*-(3-mesylpropyl)amine**

Ethereal HCl (22.5 ml, 1 M solution) was added to a stirred solution of *N*-methyl-*N*-(3-mesylpropyl)-*N*-(benzyloxycarbonyl)amine (Method 14; 5 g) in methanol (100 ml). 10%Pd/C (1.25 g) was added and the mixture was stirred under an atmosphere of hydrogen for 2 hrs. The catalyst was removed by filtration and the filtrate evaporated to dryness. Trituration with ether gave the product as a white solid. NMR ($\text{DMSO}-d_6 + d_4$ -acetic acid) 2.11 (m, 2H), 2.59 (s, 3H), 3.0 (s, 3H), 3.05 (t, 2H), 3.26 (t, 2H); m/z 152.

Method 14

***N*-Methyl-*N*-(3-mesylpropyl)-*N*-(benzyloxycarbonyl)amine**

3-chloroperoxybenzoic acid (19.2 g) was added portionwise to a stirred solution of *N*-methyl-*N*-(3-methylthiopropyl)-*N*-(benzyloxycarbonyl)amine (Method 15; 9 g) in DCM (250 ml). When the addition was completed the reaction was stirred at ambient temperature for 16 hours. The mixture was washed with sat. sodium bicarbonate solution (2 x 100 ml), brine, dried and evaporated to dryness. The residue was purified by flash column chromatography eluting with EtOAc/ iso-hexane(1:1, 3:1) and EtOAc to give the product as a solid foam. NMR (CDCl_3) 2.1 (m, 2H), 2.7-3.1 (m, 8H), 3.45 (t, 2H), 5.13 (s, 2H), 7.35 (m, 5H); m/z 286.

Method 15

***N*-Methyl-*N*-(3-methylthiopropyl)-*N*-(benzyloxycarbonyl)amine**

Sodium hydride (2.18 g, 60% in oil) was washed with iso-hexane and dried under a stream of nitrogen. DMA (50 ml) was added and the suspension was cooled to approx. 5 °C. under an atmosphere of nitrogen. A solution of *N*-(3-methylthiopropyl)-*N*-(benzyloxycarbonyl)amine (Method 16; 10 g) in DMA (150 ml)

was added dropwise keeping the temperature below 10°C. The reaction mixture was allowed to warm to ambient temperature, stirred for 30 mins. and then cooled to 0°C. Methyl iodide (8.9 g) was added dropwise and the reaction then stirred at ambient temperature for 16 hours. The solvent was evaporated under reduced pressure and the residue partitioned between

5 EtOAc and sat. ammonium chloride solution. The organic layer was separated, washed with brine and dried. Evaporation under reduced pressure gave the product as a yellow oily gum. NMR (CDCl₃) 1.83 (m, 2H), 2.07 (brs, 3H), 2.45 (m, 2H), 2.94 (s, 3H), 3.37 (t, 2H), 5.12 (s, 2H), 7.34 (m, 5H); m/z 254.

10 **Method 16**

N-(3-Methylthiopropyl)-*N*-(benzyloxycarbonyl)amine

Benzyl chloroformate (17.8 g) was added dropwise to a cooled solution of 3-methyl thiopropylamine (10 g) and triethylamine (10.6 g) in DCM (250 ml). The reaction mixture was kept at 0°C during the addition and then at ambient temperature for 16 hours. The

15 mixture was washed with aqueous citric acid (1 M, 100 ml), brine and dried. The residue was chromatographed eluting with EtOAc/iso-hexane(7:3) to give the product as a colourless gum. NMR (CDCl₃) 1.8 (m, 2H), 2.08 (s, 3H), 2.52 (t, 2H), 3.3 (m, 2H), 4.9 (brs, 1H), 5.1 (s, 2H), 7.35 (m, 5H); m/z 240.

20 **Method 17**

4-Methyl-4-hydroxy-3-methylaminotetrahydropyran.

A solution 4-methyl-3,4-epoxytetrahydropyran (Method 18; 26 g) and methylamine (33%) in ethanol (250 ml) was allowed to stand at room temperature for 5 days. The mixture was evaporated and the residue recrystallized from toluene (170 ml). M/z 145 [M⁺]; m.p.

25 (toluene) 110-112°C.

Method 18

4-Methyl-3,4-epoxytetrahydropyran

5,6-Dihydro-4-methyl-2H-pyran (Bull. Soc. Chim. Fr., 1967, 8, 2989-96; 30 g) was

30 dissolved in DCM (500 ml). *M*-chloroperoxybenzoic acid (63 g) was added over 20 minutes and the mixture was then refluxed for 4 hours. The solid was filtered and the filtrate washed

with 10% sodium sulphite, 10% sodium hydroxide, brine and then the organic layer was dried and evaporated to leave the product as an oil which was used without further purification.

Method 19

5 1-(2,3,5,6-Tetrafluoropyridin-4-yl)piperazine hydrochloride

To an ice bath cooled, stirred solution of 1-(2,3,5,6-tetrafluoropyridin-4-yl)-4-benzylpiperazine (Method 20; 4.59 g, 14.1 mmol) in 1,2-dichloroethane (40 ml) was added chloroethyl chloroformate (1.6 ml, 14.7 mmol) over five minutes. The solution was heated to reflux and stirred for two hours and allowed to cool
10 to ambient temperature. Volatile material was removed by evaporation. The residue was dissolved in methanol, heated to reflux and stirred for 2.5 hours, allowed to cool to ambient temperature. Volatile material was removed by evaporation. The residue was purified by chromatography eluting with 5-10% MeOH in DCM to give a solid which was triturated with ether to give the title compound (2.57 g) as a solid. NMR 3.05 (m, 4H), 3.53 (m, 4H); m/z
15 236.

Method 20

1-(2,3,5,6-Tetrafluoropyridin-4-yl)-4-benzylpiperazine

To a solution of pentafluoropyridine (1.76 ml, 16.0 mmol) and 1-benzylpiperazine
20 (2.64 g, 15.0 mmol) in DMSO (20 ml) was added potassium carbonate (4.20 g, 30.4 ml). The suspension was heated to 100°C and stirred for five hours and allowed to cool to ambient temperature, poured onto water and then extracted with DCM. The extract was washed with brine, water, and dried. Volatile material was removed by evaporation to give the title compound (4.59 g) as an oil. NMR (CDCl₃) 2.58 (m, 4H), 3.52 (m, 4H), 3.57 (s, 2H),
25 7.24-7.38 (m, 5H and CDCl₃); m/z 326.

Method 21

3-Cyanomethylpiperidine

1-Benzylloxycarbonyl-3-cyanomethylpiperidine (Method 22; 12.5 g) was dissolved in
30 ethanol (20 ml) and ethereal HCl (1 ml) and 5% palladium on charcoal (2.0 g) was added. The mixture was shaken under an atmosphere of hydrogen for 2.5 hours. The catalyst was filtered

and the filtrate evaporated *in vacuo*. The crude product was dissolved in ethanol and recrystallized from ether. M/z 124 [M⁺].

Method 22

5 1-Benzylloxycarbonyl-3-cyanomethylpiperidine

1-Benzylloxycarbonyl-3-mesyloxymethylpiperidine (WO 0100207; 8.4 g) was dissolved in DMSO (100 ml) and sodium cyanide (1.6 g) was added. The mixture was heated to 100°C and stirred continuously under an argon atmosphere for 6 hours. The mixture was cooled and poured into water (200 ml). The product was extracted with EtOAc, washed with
10 ferrous sulphate solution (10% w/w), dried and evaporated *in vacuo*. TLC (40% EtOAc/petroleum ether (60/80) Rf. 0.54.

Method 23

1-(2-Methylaminoethyl)-1,2,4-triazole

15 To a cooled solution (ice bath) of 1-{2-[N-methyl-N-(*t*-butoxycarbonyl)amino]ethyl}-1,2,4-triazole (Method 24; 1.6 g, 7.2 mmol) in DCM (10 ml) was added trifluoroacetic acid (5 ml). The reaction mixture was stirred for 3 hours at 0°C and then at room temperature for 1 hour. The reaction mixture was evaporated to dryness with toluene and the crude product was dissolved in DCM (3 ml) and
20 concentrated HCl in ether was added dropwise under vigorous stirring to yield the title compound as a white solid. M/z 126.9.

Method 24

1-{2-[N-Methyl-N-(*t*-butoxycarbonyl)amino]ethyl}-1,2,4-triazole

25 To a cooled solution (ice bath) of 2-[N-methyl-N-(*t*-butoxycarbonyl)amino]ethanol (J. Med. Chem., 1999, 42(11), 2007-2020; 5 g, 28 mmol), triphenyl phosphine (9.35 g, 36 mmol) and 1,2,4-triazole (1.64 g, 24 mmol) in DCM (30 ml) was added diethylazodicarboxylate (5.64 ml, 35.7 mmol) dropwise. The ice bath was removed and the reaction mixture allowed to stir at room temperature for 1 hour. The precipitate was discarded and the filtrate was
30 concentrated and purified by flash chromatography, eluting with a 50% EtOAc / DCM and then 100 % EtOAc to give a pale yellow oil which was used without further purification.

Example 194

The following illustrate representative pharmaceutical dosage forms containing the compound of formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof (hereafter compound X), for therapeutic or prophylactic use in humans:-

(a): Tablet I	mg/tablet
Compound X	100
Lactose Ph.Eur	182.75
Croscarmellose sodium	12.0
Maize starch paste (5% w/v paste)	2.25
Magnesium stearate	3.0

5

(b): Tablet II	mg/tablet
Compound X	50
Lactose Ph.Eur	223.75
Croscarmellose sodium	6.0
Maize starch	15.0
Polyvinylpyrrolidone (5% w/v paste)	2.25
Magnesium stearate	3.0

(c): Tablet III	mg/tablet
Compound X	1.0
Lactose Ph.Eur	93.25
Croscarmellose sodium	4.0
Maize starch paste (5% w/v paste)	0.75
Magnesium stearate	1.0

(d): Capsule	mg/capsule
Compound X	10
Lactose Ph.Eur	488.5
Magnesium stearate	1.5

- 94 -

(e): Injection I	(50 mg/ml)
Compound X	5.0% w/v
1 M Sodium hydroxide solution	15.0% v/v
0.1 M Hydrochloric acid	(to adjust pH to 7.6)
Polyethylene glycol 400	4.5% w/v
Water for injection	to 100%

(f): Injection II	10 mg/ml
Compound X	1.0% w/v
Sodium phosphate BP	3.6% w/v
0.1 M Sodium hydroxide solution	15.0% v/v
Water for injection	to 100%


(g): Injection III	(1 mg/ml, buffered to pH6)
Compound X	0.1% w/v
Sodium phosphate BP	2.26% w/v
Citric acid	0.38% w/v
Polyethylene glycol 400	3.5% w/v
Water for injection	to 100%

5 Note

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.

Claims


1. A compound of formula (I):




(I)

wherein:

X is a group of formula (A) or (B):



(A)



(B)

- 10 **R¹** is cyano, halo, trifluoromethyl, trifluoromethoxy, C₁₋₄alkyl, C₁₋₄alkoxy, N-(C₁₋₄alkyl)amino or N,N-(C₁₋₄alkyl)₂amino;
R² is halo, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, C₁₋₄alkyl or C₁₋₄alkoxy;
A is —NR^a—, —O— or a direct bond; wherein R^a is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl; wherein R^a may be optionally substituted by one or more R⁵;
 15 **R³** is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl or C₂₋₁₀alkynyl wherein R³ may be optionally substituted by one or more R⁶; or R³ is carbocyclyl or heterocyclyl wherein R³ may be optionally substituted on carbon by one or more R⁷; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R⁸;
R⁴ is halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy,
 20 carbamoyl, mercapto, sulphamoyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy, C₁₋₄alkanoyl, C₁₋₄alkanoyloxy, N-(C₁₋₄alkyl)amino, N,N-(C₁₋₄alkyl)₂amino, C₁₋₄alkanoylamino, N-(C₁₋₄alkyl)carbamoyl, N,N-(C₁₋₄alkyl)₂carbamoyl, N-(carbocyclyl)carbamoyl, N,N-(carbocyclyl)₂carbamoyl, N-(heterocyclyl)carbamoyl, N,N-(heterocyclyl)₂carbamoyl, C₁₋₄alkylS(O)_a wherein a is 0 to 2, C₁₋₄alkoxycarbonyl,
 25 N-(C₁₋₄alkyl)sulphamoyl, N,N-(C₁₋₄alkyl)₂sulphamoyl, C₁₋₄alkylsulphonylamino or (nitrogen-linked heterocyclic ring)carbonyl;

- R^5 and R^6 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, C_{1-6} alkanoylamino, C_{2-6} alkenyloxycarbonyl, C_{1-6} alkoxycarbonyl, heterocyclyl, heterocyclyloxy, heterocyclylamino,
- 5 heterocyclyl- N -(C_{1-6} alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl, heterocyclylcarbonylamino, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylamino, carbocyclyl- N -(C_{1-6} alkyl)amino, carbocyclylsulphonyl, carbocyclylcarbonyl, carbocyclylcarbonylamino, carbocyclyloxycarbonyl, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkoxycarbonylamino, N -(C_{1-6} alkyl)carbamoyl,
- 10 N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkylS(O)_a wherein a is 0-2, N -(C_{1-6} alkyl)sulphamoylamino, N,N -(C_{1-6} alkyl)₂sulphamoylamino, C_{1-6} alkylsulphonylamino, (C_{1-6} alkyl)sulphonyl- N -(C_{1-6} alkyl)amino, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl)₂sulphamoyl; wherein R^5 and R^6 may be optionally substituted on carbon by one or more R^9 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may
- 15 be independently optionally substituted by R^{10} ;

- R^7 and R^9 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, trifluoromethyl, trifluoromethoxy, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, C_{1-4} alkoxycarbonylamino, C_{2-4} alkenyloxycarbonyl, C_{1-4} alkanoyl, C_{1-4} alkanoylamino,
- 20 C_{1-4} alkanoyloxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0-2, N -(C_{1-4} alkyl)sulphamoylamino, N,N -(C_{1-4} alkyl)₂sulphamoylamino, (C_{1-4} alkyl)sulphonylamino, (C_{1-4} alkyl)sulphonyl- N -(C_{1-4} alkyl)amino, N -(C_{1-4} alkyl)sulphamoyl, N,N -(C_{1-4} alkyl)₂sulphamoyl, heterocyclyl, heterocyclyloxy, heterocyclylamino,
- 25 heterocyclyl- N -(C_{1-4} alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl, heterocyclylcarbonylamino, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylamino, carbocyclyl- N -(C_{1-4} alkyl)amino, carbocyclylsulphonyl, carbocyclylcarbonyl, carbocyclylcarbonylamino, and carbocyclyloxycarbonyl; wherein R^7 and R^9 may be independently optionally substituted on carbon by one or more R^{11} ;
- 30 R^8 and R^{10} are independently selected from C_{1-4} alkyl, C_{1-4} alkanoyl, sulphamoyl, C_{1-4} alkylsulphonyl, C_{1-4} alkoxycarbonyl, carbamoyl, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, N -(C_{1-4} alkyl)sulphamoyl, N,N -(C_{1-4} alkyl)₂sulphamoyl,

heterocyclyl, heterocyclylcarbonyl, heterocyclioxycarbonyl, heterocyclsulphonyl, carbocyclyl, carbocyclylcarbonyl, carbocyclioxycarbonyl and carbocyclsulphonyl; wherein R^8 and R^{10} may be independently optionally substituted on carbon by one or more R^{12} ;

- R^{11} and R^{12} are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, vinyl, allyl, methoxy, ethoxy, vinyloxy, allyloxy, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, *N*-methylcarbamoyl, *N,N*-dimethylcarbamoyl, methylthio, methylsulphinyl, mesyl, *N*-methylsulphamoyl, *N,N*-dimethylsulphamoyl, heterocyclyl, heterocyclloxy, heterocyclylcarbonyl, heterocyclloxycarbonyl, carbocyclyl, carbocyclloxy, carbocyclylcarbonyl and carbocyclloxycarbonyl;
- x is 0 - 4; wherein the values of R^1 may be the same or different;
- y is 0 - 3; wherein the values of R^2 may be the same or different;
- z is 0 - 3; wherein the values of R^4 may be the same or different; and
- n is 0 - 2;

- or a pharmaceutically acceptable salt, prodrug or solvate thereof; with the provisos when X is a group of formula (A):
- 1) when R^3 is a nitrogen linked heterocyclyl, A is a direct bond;
 - 2) when x and y are 0 and n is 2, the group $R^3-A-C(O)-NH-$ is not 2-formamido, 2-acetamido, 3-acetamido, 2-propionamido, 3-[2-(fur-2-ylcarbonylmethyl)acetamido], 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido), 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido), 2-(2-phthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido), 2-(2-amino-2-benzylacetamido), 3-(2-phthalimidoacetamido), 3-(3-phthalimidopropionamido), 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido), 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido), 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-benzylacetamido) or 2-(3,3,3-trifluoro-2-hydroxy-2-methylpropionamido);
 - 3) when x and y are 0 and n is 1, the group $R^3-A-C(O)-NH-$ is not 3-acetamido, 3-(2-phthalimidoacetamido), 3-(3-phthalimidopropionamido), 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido), 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido),

- 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-isobutylacetamido) or
 3-(2-amino-2-benzylacetamido);
- 4) when x and y are 0 and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-benzamido, 2-acetamido, 2-benzyloxycarbonylamino, 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido),
- 5 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido),
 2-(2-phthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido),
 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido),
 2-(2-amino-2-benzylacetamido), 3-(trifluoroacetamido), 3-benzyloxycarbonylamino or
 4-t-butyloxycarbonylamino;
- 10 5) when $(R^1)_x$ is 7-fluoro, y is 0 and n is 2, the group $R^3-A-C(O)-NH-$ is not 3-acetamido;
- 6) when x is 0, $(R^2)_y$ is 1-cyano and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido;
- 7) when x is 0, $(R^2)_y$ is 3-bromo and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido or
 2-benzamido;
- 8) when x is 0, $(R^2)_y$ is 1-bromo and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-acetamido or
- 15 2-benzamido; and
- 9) when x is 0, $(R^2)_y$ is 1-chloro or 4-chloro and n is 0, the group $R^3-A-C(O)-NH-$ is not
 2-acetamido.

2. A compound of formula (I) as claimed in claim 1 wherein X is a group of formula (A)
 20 or a pharmaceutically acceptable salt, prodrug or solvate thereof.

3. A compound of formula (I) as claimed in claim 1 wherein X is a group of formula (B)
 or a pharmaceutically acceptable salt, prodrug or solvate thereof.

25 4. A compound of formula (I) as claimed in any one of claims 1 - 3 wherein X is in the
 2-position of the dibenzothiophene ring or a pharmaceutically acceptable salt, prodrug or
 solvate thereof.

5. A compound of formula (I) as claimed in any one of claims 1 - 4 wherein R^2 is halo,
 30 cyano or C_{1-4} alkyl or a pharmaceutically acceptable salt, prodrug or solvate thereof.

6. A compound of formula (I) as claimed in any one of claims 1 - 5 wherein A is —NR^a—, —O— or a direct bond; wherein R^a is hydrogen, C₁₋₁₀alkyl or C₂₋₁₀alkenyl; wherein R^a may be optionally substituted by one or more R⁵; wherein

R³ is C₁₋₁₀alkyl or C₂₋₁₀alkenyl wherein R³ may be optionally substituted by one or more R⁶; or R³ is carbocyclyl or heterocyclyl wherein R³ may be optionally substituted on carbon by one or more R⁷; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R⁸;

R⁵ and R⁶ are independently selected from hydroxy, cyano, amino, C₁₋₆alkoxy, heterocyclyl, heterocyclxyloxy, carbocyclyl, carbocyclxyloxy, carbocyclyl-*N*-(C₁₋₆alkyl)amino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkoxycarbonylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, *N,N*-(C₁₋₆alkyl)₂sulphamoylamino and C₁₋₆alkylsulphonylamino; wherein R⁵ and R⁶ may be independently optionally substituted on carbon by one or more R⁹;

R⁷ and R⁹ are independently selected from halo, hydroxy, nitro, carboxy, carbamoyl, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxycarbonyl, C₁₋₄alkanoyl, C₁₋₄alkanoylamino, *N,N*-(C₁₋₄alkyl)₂amino, *N,N*-(C₁₋₄alkyl)₂carbamoyl, heterocyclyl, heterocyclylcarbonyl, carbocyclyl and carbocyclylcarbonylamino; wherein R⁷ and R⁹ may be independently optionally substituted on carbon by one or more R¹¹;

R⁸ is selected from C₁₋₄alkyl, C₁₋₄alkylsulphonyl, *N,N*-(C₁₋₄alkyl)₂sulphamoyl, heterocyclyl and carbocyclyl; wherein R⁸ may be optionally substituted on carbon by one or more R¹²;

R¹¹ and R¹² are independently selected from halo, hydroxy, cyano, carbamoyl, methyl, methoxy, allyloxy, heterocyclyl and carbocyclyl; or a pharmaceutically acceptable salt, prodrug or solvate thereof.

7. A compound of formula (I) as claimed in any one of claims 1 - 6 wherein the group R³-A- is methyl, 2-oxo-pyrrolidin-1-ylmethyl, 1,2,4-triazol-1-ylmethyl, 1,1-dioxotetrahydrothien-3-ylmethyl, 2-oxooxazolidin-3-ylmethyl, pyrid-3-yloxymethyl, 1,1-dioxothiomorpholinomethyl, cyanomethyl, 2-oxo-1,2-dihydropyrid-1-ylmethyl, 2-oxocyclopentylmethyl, succinimidomethyl, 3-benzyl-2-oxopyrrolidin-1-ylmethyl, 3-hydroxypyridazin-6-yloxymethyl, 2-pyrid-4-ylethyl, 2-methoxyethyl, 1-phenoxyethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-imidazol-1-ylethyl, succinimidoethyl, prop-2-yl, 3,3,3-trifluoro-2-hydroxyprop-2-yl, 1-morpholinoprop-2-yl, 1-pyrid-4-ylprop-2-yl,

- 2-aminoprop-2-yl, 2-(*t*-butoxycarbonylamino)prop-2-yl, 1-(1,2,4-triazol-1-yl)prop-2-yl,
2-pyrid-4-ylpropyl, *t*-butyl, 1,1,1-trifluorobut-3-yl, 1-hydroxyhex-2-yl, cyclopropyl,
3-hydroxybicyclo[2.2.1]hept-2-yl, 4-nitrophenyl, morpholino, 4-methylpiperazin-1-yl,
tetrahydropyran-4-yl, 4-hydroxymethylpiperidin-1-yl, 1-methyl-2-oxopyrrolidin-4-yl,
5 2-(pyrrolidin-1-ylmethyl)pyrrolidinyl, 3-carbamoylpiperidin-1-yl, 3-hydroxyazetidin-1-yl,
2-(allyloxymethyl)morpholino, 4-(1,4-dihydrooxazin-2-one-3-yl)piperidin-1-yl,
4-(*N,N*-dimethylsulphamoyl)piperazin-1-yl, 4-hydroxyethylpiperidin-1-yl,
4-(tetrahydrofur-2-ylmethyl)piperazin-1-yl, 4-(3-methoxypropyl)piperazin-1-yl,
4-pyrid-4-ylpiperidin-1-yl, 4-pyrid-2-ylpiperazin-1-yl, 3-(*N,N*-dimethylamino)pyrrolidin-1-yl,
10 4-carboxypiperidin-1-yl, 1-methyl-2-oxo-5-phenylpyrrolidin-4-yl,
2-oxo-5,5-dimethyltetrahydrofur-4-yl, tetrahydrofur-4-yl, 2,2-dimethyltetrahydropyran-4-yl,
1-benzyl-2-oxopyrrolidin-4-yl, 2-oxo-5-phenyltetrahydrofuryl,
2-(3-hydroxypropyl)piperidin-1-yl, 4-(2-carbamylethyl)piperazin-1-yl,
3-oxo-4-(2-methoxyethyl)piperazin-1-yl, 4-(*N,N*-dimethylamino)-4-carbamoylpiperidin-1-yl,
15 4-(2-morpholinoethyl)piperazin-1-yl, 1,4-dioxo-8-azaspiro[4.5]decan-8-yl,
1,1-dioxotetrahydrothien-3-yl, 4-ethylsulphonylpiperazin-1-yl,
4-(thieno[2,3-*d*]pyrimidin-4-yl)piperazin-1-yl, 4-(2-hydroxyethyl)piperazin-1-yl,
4-carbamoylpiperidin-1-yl, 3-methyl-3-phenylpiperidin-1-yl,
2-benzyloxycarbonylpiperidin-1-yl, 4-(*N,N*-dimethylcarbamoyl)piperidin-1-yl,
20 3-(pyrid-4-yl)pyrrolidin-1-yl, 3-(pyrid-3-yl)pyrrolidin-1-yl, 4,4-difluoropiperidin-1-yl,
4-(2-methylpyrimidin-4-yl)piperazin-1-yl, 4-(2,3,5,6-tetrafluoropyrid-4-yl)piperazin-1-yl,
4-(pyrimidin-4-yl)piperazin-1-yl, 3-cyanomethylpiperidin-1-yl,
4-cyclohexylcarbonylaminopiperidin-1-yl, 4-acetamidopiperidin-1-yl,
4-(6-chloropyrimidin-4-yl)piperazin-1-yl, 4-(pent-3-yl)piperazin-1-yl,
25 1,2,5,6-tetrahydropyrid-1-yl, 1-methylpiperidin-1-yl, 2-methylpiperidin-1-yl,
decahydroquinolin-1-yl, 3-ethoxycarbonyl-4-oxopiperidin-1-yl,
2-ethoxycarbonylpyrrolidin-1-yl, 4-acetylpiperidin-1-yl, 2-azabicyclo[2.2.1]hept-2-yl,
1,2,3,4,-tetrahydro-9H-pyrido[3,4-*b*]indol-2-yl,
4-(2,3-dihydro-2-oxobenzimidazol-1-yl)piperidin-1-yl, 4-(2-methoxyphenyl)piperazin-1-yl,
30 4-methylpiperidin-1-yl, 4-(3-chlorophenyl)piperazin-1-yl,
4-(3-methoxyphenyl)piperazin-1-yl, 4-(4-methoxyphenyl)piperazin-1-yl,
4-(4-chlorophenyl)piperazin-1-yl,

- 4-(2,3-dihydro-2-oxobenzimidazol-1-yl)-1,2,5,6-tetrahydropyrid-1-yl, 4-phenylpiperidin-1-yl,
 4-(3-fur-2-ylpyrazol-5-yl)piperidin-1-yl, 2-pyrid-4-ylethylamino,
 3-imidazol-1-ylpropylamino, 4-hydroxycyclohexylamino,
 2-(*N,N*-dimethylsulphamoylamino)ethylamino, 2-(isopropylsulphonylamino)ethylamino,
 5 2-imidazol-5-ylethylamino, 2-mesyethylamino, 2-morpholinoethylamino,
 1-methoxycarbonylcyclopropylamino, 1-benzylpyrrolidin-3-ylamino,
 3-(*N*-methylanilino)propylamino, 2-(5-methyl-2,4-dioxothiazolidin-3-yl)ethylamino,
 2-(*t*-butoxycarbonylamino)ethylamino, *N*-(*N*-methyl-*N*-pyrid-3-ylmethylaminopropyl)amino,
 1-cyclohexylethylamino, *N*-methyl-*N*-(2-pyrid-4-ylethyl)amino,
 10 *N*-methyl-*N*-(2-pyrid-2-ylethyl)amino, *N*-methyl-*N*-(2-cyanoethyl)amino,
N-methyl-*N*-(pyrid-3-ylmethyl)amino, *N*-methyl-*N*-(2-*N,N*-dimethylaminoethyl)amino,
N-methyl-*N*-(1-methylpiperidin-4-yl)amino, *N*-methyl-*N*-(3-mesylpropyl)amino,
N-methyl-*N*-(4-hydroxy-4-methyltetrahydropyran-3-yl)amino,
N-(pyrid-3-ylmethyl)-*N*-(2-cyanoethyl)amino, *N*-methyl-*N*-(2-hydroxypropyl)amino,
 15 *N*-methyl-*N*-(2,2-dimethoxyethyl)amino, *N*-methyl-*N*-phenethylamino,
N-methyl-*N*-(tetrahydrofur-2-ylmethyl)amino, *N*-methyl-*N*-(2-morpholinoethyl)amino,
N-methyl-*N*-(6-methylpyrid-2-ylmethyl)amino, *N*-methyl-*N*-(1-methylpyrrolidin-3-yl)amino,
N-methyl-*N*-[2-(4-hydroxyphenyl)-2-hydroxyethyl]amino,
N-methyl-*N*-(1-benzylpyrrolidin-3-yl)amino, *N*-methyl-*N*-[2-(1,2,4-triazol-1-yl)ethyl]amino,
 20 *N*-methyl-*N*-(fur-2-ylmethyl)amino, *N*-methyl-*N*-(benzimidazol-2-ylmethyl)amino,
N-methyl-*N*-benzylamino, *N*-methyl-*N*-(2-chlorobenzyl)amino,
N-methyl-*N*-(3-chlorobenzyl)amino, *N*-methyl-*N*-(4-chlorobenzyl)amino,
N-methyl-*N*-[2-(3,4-dimethoxypyrid-4-yl)ethyl]amino,
N-methyl-*N*-(5-phenylpyrazol-3-ylmethyl)amino, *N*-methyl-*N*-(4-fluorobenzyl)amino,
 25 *N*-methyl-*N*-(2-methoxyphenylprop-2-yl)amino, *N*-ethyl-*N*-(pyrid-4-ylmethyl)amino,
N-(2-methoxyethyl)-*N*-(pyrid-3-ylmethyl)amino, *N*-ethyl-*N*-(2-methoxyethyl)amino,
N-(2-hydroxyethyl)-*N*-isopropylamino, *N*-(2-cyanoethyl)-*N*-(3-morpholinopropyl)amino,
N-(2-cyanoethyl)-*N*-(thien-2-ylmethyl)amino, *N*-(2-cyanoethyl)-*N*-benzylamino,
N-ethyl-*N*-(1-benzylpyrrolidin-3-yl)amino, *N*-ethyl-*N*-(1,1-dioxotetrahydrothien-3-yl)amino,
 30 *N*-(2-pyrid-4-ylethyl)-*N*-cyclopropylamino,
N-(2-hydroxy-2-pyrid-4-ylethyl)-*N*-isopropylamino,

N-(4-hydroxycyclohexyl)-*N*-isopropylamino, *N*-allyl-*N*-(1,1-dioxotetrahydrothien-3-yl)amino, diallylamino, *N*-allyl-*N*-cyclopentylamino or *N*-benzyl-*N*-propylamino.

8. A compound of formula (I) as claimed in any one of claims 1 - 7 wherein R⁴ is
- 5 C₁₋₄alkyl or a pharmaceutically acceptable salt, prodrug or solvate thereof.
9. A compound of formula (I) as claimed in any one of claims 1 - 8 wherein x is 0 or a pharmaceutically acceptable salt, prodrug or solvate thereof.
- 10 10. A compound of formula (I) as claimed in any one of claims 1 - 9 wherein y is 0-1 or a pharmaceutically acceptable salt, prodrug or solvate thereof.
11. A compound of formula (I) as claimed in any one of claims 1 - 10 wherein z is 1 or a pharmaceutically acceptable salt, prodrug or solvate thereof.
- 15 12. A compound of formula (I) as claimed in any one of claims 1 - 11 wherein n is 2 or a pharmaceutically acceptable salt, prodrug or solvate thereof.
13. A compound of formula (I) (as depicted in claim 1) wherein:
- 20 X is a group of formula (A) (as depicted in claim 1) or a group of formula (B) (as depicted in claim 1) in the 2-position of the dibenzothiophene ring;
R² is bromo, cyano or methyl;
A is —NR^a—, —O— or a direct bond; wherein R^a is hydrogen, methyl, ethyl, propyl, isopropyl or allyl; wherein said methyl, ethyl, propyl, isopropyl or allyl may be optionally
- 25 substituted by one or more R⁵;
R³ is methyl, ethyl, propyl, butyl, hexyl or allyl wherein R³ may be optionally substituted by one or more R⁶; or R³ is selected from cyclopropyl, cyclopentyl, cyclohexyl, bicyclo[2,2,1]heptyl, phenyl, tetrahydropyranyl, morpholino, 2-oxopyrrolidinyl, piperidinyl, pyrrolidinyl, azetidiny, tetrahydrofuryl, 1,4-dioxo-8-azaspiro[4.5]decanyl,
- 30 1,1-dioxotetrahydrothienyl, 1,2,3,4,-tetrahydro-9H-pyrido[3,4-b]indole, decahydroquinolin-1-yl, 1,2,5,6-tetrahydropyridyl or piperazinyl; wherein R³ may be

optionally substituted on carbon by one or more R⁷; and wherein if any heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R⁸;

R⁵ is selected from cyano, methoxy, pyridyl;

- R⁶ is selected from hydroxy, cyano, amino, methoxy, pyridyl, 2-oxopyrrolidinyl,
 5 1,2,4-triazolyl, 1,1-dioxotetrahydrothienyl, thienyl, 2-oxooxazolidinyl, imidazolyl,
 1,1-dioxothiomorpholino, 2-oxo-1,2-dihydropyridyl, benzimidazolyl, pyrazolyl, succinimido,
 tetrahydrofuryl, 2,4-dioxothiazolidinyl, morpholino, furyl, pyridyloxy, pyridazinyloxy,
 cyclopentyl, cyclohexyl, phenyl, phenoxy, *N*-methylanilino, *N,N*-dimethylamino,
t-butoxycarbonylamino, mesyl, *N,N*-dimethylsulphamoylamino and
 10 isopropylsulphonylamino; wherein R⁶ may be optionally substituted on carbon by one or more
 R⁹;

- R⁷ is selected from fluoro, hydroxy, nitro, carboxy, carbamoyl, methyl, ethyl, propyl,
 methoxycarbonyl, ethoxycarbonyl, acetyl, acetamido, *N,N*-dimethylamino,
N,N-dimethylcarbamoyl, *N,N*-diethylcarbamoyl, 1,4-dihydrooxazin-2-one, pyrazolyl,
 15 2-oxo-2,3-dihydrobenzimidazolyl, pyridyl, 2,3-dihydro-2-oxobenzimidazolyl,
 morpholinocarbonyl, phenyl and cyclohexylcarbonylamino; wherein any R⁷ may be optionally
 substituted on carbon by one or more R¹¹;

R⁹ is selected from fluoro, chloro, hydroxy, methyl, methoxy, pyridyl and phenyl;

- R⁸ is selected from methyl, ethyl, propyl, pentyl, ethylsulphonyl,
 20 *N,N*-dimethylsulphamoyl, thieno[2,3-*d*]pyrimidinyl, pyrimidinyl, pyridyl and phenyl; wherein
 any R⁸ may be optionally substituted on carbon by one or more R¹²;

R¹¹ is selected from hydroxy, cyano, allyloxy, pyrrolidinyl, furyl and phenyl;

R¹² is selected from fluoro, chloro, hydroxy, carbamoyl, methyl, methoxy,
 tetrahydrofuryl, morpholino and phenyl;

- 25 R⁴ is methyl;

R⁴ is C₁₋₄alkyl;

x is 0;

y is 0-1;

z is 1; and

- 30 n is 0-2;

or a pharmaceutically acceptable salt, prodrug or solvate thereof;

with the provisos when X is a group of formula (A):

- 104 -

- 1) when R³ is a nitrogen linked heterocyclyl, A is a direct bond;
- 2) when x and y are 0 and n is 2, the group R³-A-C(O)-NH- is not 2-acetamido, 2-propionamido, 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido) or 2-(2-amino-2-benzylacetamido);
- 5 3) when x and y are 0 and n is 0, the group R³-A-C(O)-NH- is not 2-benzamido, 2-acetamido, 2-benzoyloxycarbonylamino, 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido) or 2-(2-amino-2-benzylacetamido);
- 4) when x is 0, (R²)_y is 1-cyano and n is 0, the group R³-A-C(O)-NH- is not 2-acetamido;
- 5) when x is 0, (R²)_y is 3-bromo and n is 0, the group R³-A-C(O)-NH- is not 2-acetamido or
- 10 2-benzamido; and
- 6) when x is 0, (R²)_y is 1-bromo and n is 0, the group R³-A-C(O)-NH- is not 2-acetamido or 2-benzamido.


14. A compound of formula (I) selected from:

- 15 2-(N'-pyrid-4-ylethyl-N'-methylureido)dibenzothiophene;
- 5,5-dioxo-2-(N'-pyrid-4-ylethyl-N'-methylureido)dibenzothiophene;
- 5,5-dioxo-2-[4-(pyrid-4-yl)piperidin-1-ylcarbonylamino]dibenzothiophene;
- 5,5-dioxo-2-[4-(pyrid-2-yl)piperidin-1-ylcarbonylamino]dibenzothiophene;
- 5,5-dioxo-2-(N'-phenethyl-N'-methylureido)dibenzothiophene;
- 20 5,5-dioxo-2-(4,4,4-trifluoro-2-methylbutyrylamino)dibenzothiophene;
- 5,5-dioxo-2-[4-(2,3,5,6-tetrafluoropyrid-4-yl)piperazin-1-ylcarbonylamino]dibenzothiophene;
- or a pharmaceutically acceptable salt, prodrug or solvate thereof.

15. A process for preparing a compound of formula (I) as claimed in any one of claims
- 25 1-14 or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof which process (wherein R¹, R², R³, A and n are, unless otherwise specified, as defined in formula (I)) comprises of:


Process a): for compounds of formula (I) wherein A is a direct bond and X is a group of formula (A); reacting an amine of formula (II):

- 105 -



(II)

with an acid of formula (III):



(III)

5


or an activated derivative thereof; or

Process b): for compounds of formula (I) wherein $n > 0$; by oxidising a compound of formula

(I) where $n = 0$;

Process c): for compounds of formula (I) wherein A is $-NR^a-$ and X is a group of formula (A);

10 by reacting a compound of formula (IV):



(IV)


wherein L is a displaceable group; with an amine of formula (V):



(V)

15

Process d): for compounds of formula (I) wherein A is $-NR^a-$ or $-O-$ and X is a group of formula (A); reacting a compound of formula (II) with a compound of formula (VI):



(VI)


20 *Process e)*: for compounds of formula (I) wherein A is $-NH-$ and X is a group of formula (A); reacting a compound of formula (II) with an isocyanate of formula (VII):

- 106 -



(VII)

Process f): for compounds of formula (I) wherein X is a group of formula (B) reacting a compound of formula (II) with a compound of formula (VIII):



(VIII)

wherein L is a displaceable group;


and thereafter if necessary:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

16. A pharmaceutical composition which comprises a compound of the formula (I) or a pharmaceutically acceptable salt, prodrug or solvate thereof as claimed in any one of claims

1-14, in association with a pharmaceutically acceptable diluent or carrier.


17. A compound of formula (IA'):




(IA')

20 wherein:

X is a group of formula (A) or (B):



(A)



(B)

R^1 is cyano, halo, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl, C_{1-4} alkoxy, $N-(C_{1-4}alkyl)amino$ or $N,N-(C_{1-4}alkyl)_2amino$;

R^2 is halo, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, C_{1-4} alkyl or C_{1-4} alkoxy;

A is $—NR^a—$, $—O—$ or a direct bond; wherein R^a is hydrogen, $C_{1-10}alkyl$,

5 $C_{2-10}alkenyl$, $C_{2-10}alkynyl$; wherein R^a may be optionally substituted by one or more R^5 ;

R^3 is hydrogen, $C_{1-10}alkyl$, $C_{2-10}alkenyl$ or $C_{2-10}alkynyl$ wherein R^3 may be optionally substituted by one or more R^6 ; or R^3 is carbocyclyl or heterocyclyl wherein R^3 may be optionally substituted on carbon by one or more R^7 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by R^8 ;

10 R^4 is halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, $C_{1-4}alkoxycarbonyl$, $C_{1-4}alkyl$, $C_{2-4}alkenyl$, $C_{2-4}alkynyl$, $C_{1-4}alkoxy$, $C_{1-4}alkanoyl$, $C_{1-4}alkanoyloxy$, $N-(C_{1-4}alkyl)amino$, $N,N-(C_{1-4}alkyl)_2amino$, $C_{1-4}alkanoylamino$, $N-(C_{1-4}alkyl)carbamoyl$, $N,N-(C_{1-4}alkyl)_2carbamoyl$, $N-(carbocyclyl)carbamoyl$, $N,N-(carbocyclyl)_2carbamoyl$, $N-(heterocyclyl)carbamoyl$,
15 $N,N-(heterocyclyl)_2carbamoyl$, $C_{1-4}alkylS(O)_a$ wherein a is 0 to 2, $C_{1-4}alkoxycarbonyl$, $N-(C_{1-4}alkyl)sulphamoyl$, $N,N-(C_{1-4}alkyl)_2sulphamoyl$, $C_{1-4}alkylsulphonylamino$ or (nitrogen-linked heterocyclic ring)carbonyl;

R^5 and R^6 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, $C_{1-6}alkoxy$,

20 $C_{1-6}alkanoyl$, $C_{1-6}alkanoyloxy$, $C_{1-6}alkanoylamino$, $C_{2-6}alkenyloxycarbonyl$, $C_{1-6}alkoxycarbonyl$, heterocyclyl, heterocycliloxy, heterocyclylamino, heterocyclyl- $N-(C_{1-6}alkyl)amino$, heterocyclylsulphonyl, heterocyclylcarbonyl, heterocyclylcarbonylamino, heterocycliloxycarbonyl, carbocyclyl, carbocycliloxy, carbocyclylamino, carbocyclyl- $N-(C_{1-6}alkyl)amino$, carbocyclylsulphonyl,
25 carbocyclylcarbonyl, carbocyclylcarbonylamino, carbocycliloxycarbonyl, $N-(C_{1-6}alkyl)amino$, $N,N-(C_{1-6}alkyl)_2amino$, $C_{1-6}alkoxycarbonylamino$, $N-(C_{1-6}alkyl)carbamoyl$, $N,N-(C_{1-6}alkyl)_2carbamoyl$, $C_{1-6}alkylS(O)_a$ wherein a is 0-2, $N-(C_{1-6}alkyl)sulphamoylamino$, $N,N-(C_{1-6}alkyl)_2sulphamoylamino$, $C_{1-6}alkylsulphonylamino$, $(C_{1-6}alkyl)sulphonyl-N-(C_{1-6}alkyl)amino$, $N-(C_{1-6}alkyl)sulphamoyl$ and

30 $N,N-(C_{1-6}alkyl)_2sulphamoyl$; wherein R^5 and R^6 may be optionally substituted on carbon by one or more R^9 ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be independently optionally substituted by R^{10} ;

- R^7 and R^9 are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, trifluoromethyl, trifluoromethoxy, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, C_{1-4} alkoxycarbonylamino, C_{2-4} alkenyloxycarbonyl, C_{1-4} alkanoyl, C_{1-4} alkanoylamino, 5 C_{1-4} alkanoyloxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0-2, N -(C_{1-4} alkyl)sulphamoylamino, N,N -(C_{1-4} alkyl)₂sulphamoylamino, (C_{1-4} alkyl)sulphonylamino, (C_{1-4} alkyl)sulphonyl- N -(C_{1-4} alkyl)amino, N -(C_{1-4} alkyl)sulphamoyl, N,N -(C_{1-4} alkyl)₂sulphamoyl, heterocyclyl, heterocyclyloxy, heterocyclylamino, 10 heterocyclyl- N -(C_{1-4} alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl, heterocyclylcarbonylamino, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylamino, carbocyclyl- N -(C_{1-4} alkyl)amino, carbocyclylsulphonyl, carbocyclylcarbonyl, carbocyclylcarbonylamino, and carbocyclyloxycarbonyl; wherein R^7 and R^9 may be independently optionally substituted on carbon by one or more R^{11} ;
- 15 R^8 and R^{10} are independently selected from C_{1-4} alkyl, C_{1-4} alkanoyl, sulphamoyl, C_{1-4} alkylsulphonyl, C_{1-4} alkoxycarbonyl, carbamoyl, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, N -(C_{1-4} alkyl)sulphamoyl, N,N -(C_{1-4} alkyl)₂sulphamoyl, heterocyclyl, heterocyclylcarbonyl, heterocyclyloxycarbonyl, heterocyclylsulphonyl, carbocyclyl, carbocyclylcarbonyl, carbocyclyloxycarbonyl and carbocyclylsulphonyl; wherein 20 R^8 and R^{10} may be independently optionally substituted on carbon by one or more R^{12} ;
- R^{11} and R^{12} are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, vinyl, allyl, methoxy, ethoxy, vinyloxy, allyloxy, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, N -methylcarbamoyl, N,N -dimethylcarbamoyl, 25 methylthio, methylsulphinyl, mesyl, N -methylsulphamoyl, N,N -dimethylsulphamoyl, heterocyclyl, heterocyclyloxy, heterocyclylcarbonyl, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylcarbonyl and carbocyclyloxycarbonyl;
- x is 0 - 4; wherein the values of R^1 may be the same or different;
- y is 0 - 3; wherein the values of R^2 may be the same or different;
- 30 z is 0 - 3; wherein the values of R^4 may be the same or different; and
- n is 0 - 2;
- or a pharmaceutically acceptable salt, prodrug or solvate thereof;

- 109 -


with the provisos when X is a group of formula (A):

- 1) when R^3 is a nitrogen linked heterocyclyl, A is a direct bond;
- 2) when x and y are 0 and n is 2, the group $R^3-A-C(O)-NH-$ is not 2-formamido, 2-acetamido, 2-propionamido, 2-(2-phthalimidoacetamido), 2-(3-phthalimidopropionamido),
 5 2-(2-phthalimido-2-isopropylacetamido), 2-(2-phthalimido-2-isobutylacetamido),
 2-(2-phthalimido-2-benzylacetamido), 2-(2-aminoacetamido), 2-(3-aminopropionamido),
 2-(2-amino-2-isopropylacetamido), 2-(2-amino-2-isobutylacetamido),
 2-(2-amino-2-benzylacetamido), 3-(2-phthalimidoacetamido), 3-(3-phthalimidopropionamido),
 3-(2-phthalimido-2-isopropylacetamido), 3-(2-phthalimido-2-isobutylacetamido),
 10 3-(2-phthalimido-2-benzylacetamido), 3-(2-aminoacetamido), 3-(3-aminopropionamido),
 3-(2-amino-2-isopropylacetamido), 3-(2-amino-2-benzylacetamido) or
 2-(3,3,3-trifluoro-2-hydroxy-2-methylpropionamido);
- 3) when x and y are 0 and n is 1, the group $R^3-A-C(O)-NH-$ is not 3-(2-phthalimidoacetamido),
 3-(3-phthalimidopropionamido), 3-(2-phthalimido-2-isopropylacetamido),
 15 3-(2-phthalimido-2-isobutylacetamido), 3-(2-phthalimido-2-benzylacetamido),
 3-(2-aminoacetamido), 3-(3-aminopropionamido), 3-(2-amino-2-isopropylacetamido),
 3-(2-amino-2-isobutylacetamido) or 3-(2-amino-2-benzylacetamido);
- 4) when x and y are 0 and n is 0, the group $R^3-A-C(O)-NH-$ is not 2-(2-phthalimidoacetamido),
 2-(3-phthalimidopropionamido), 2-(2-phthalimido-2-isopropylacetamido),
 20 2-(2-phthalimido-2-isobutylacetamido), 2-(2-phthalimido-2-benzylacetamido),
 2-(2-aminoacetamido), 2-(3-aminopropionamido), 2-(2-amino-2-isopropylacetamido),
 2-(2-amino-2-isobutylacetamido) or 2-(2-amino-2-benzylacetamido);
- 5) when $(R^1)_x$ is 7-fluoro, y is 0 and n is 2, the group $R^3-A-C(O)-NH-$ is not 3-acetamido;
 or a pharmaceutically acceptable salt, prodrug or solvate thereof, for use as a medicament.

25



18. A method of treating disorders mediated by the neuropeptide Y5 receptor in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB'):

- 110 -



wherein:

X is a group of formula **(A)** or **(B)**:



R¹ is cyano, halo, trifluoromethyl, trifluoromethoxy, C₁₋₄alkyl, C₁₋₄alkoxy,

N-(C₁₋₄alkyl)amino or *N,N*-(C₁₋₄alkyl)₂amino;

R² is halo, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, C₁₋₄alkyl or C₁₋₄alkoxy;

A is —NR^a—, —O— or a direct bond; wherein R^a is hydrogen, C₁₋₁₀alkyl,

C₂₋₁₀alkenyl, C₂₋₁₀alkynyl; wherein R^a may be optionally substituted by one or more R⁵;

R³ is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl or C₂₋₁₀alkynyl wherein R³ may be optionally substituted by one or more R⁶; or R³ is carbocyclyl or heterocyclyl wherein R³ may be optionally substituted on carbon by one or more R⁷; and wherein if said heterocyclyl contains

an -NH- moiety that nitrogen may be optionally substituted by R⁸;

R⁴ is halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl,

C₁₋₄alkoxy, C₁₋₄alkanoyl, C₁₋₄alkanoyloxy, *N*-(C₁₋₄alkyl)amino, *N,N*-(C₁₋₄alkyl)₂amino,

C₁₋₄alkanoylamino, *N*-(C₁₋₄alkyl)carbamoyl, *N,N*-(C₁₋₄alkyl)₂carbamoyl,

N-(carbocyclyl)carbamoyl, *N,N*-(carbocyclyl)₂carbamoyl, *N*-(heterocyclyl)carbamoyl,

N,N-(heterocyclyl)₂carbamoyl, C₁₋₄alkylS(O)_a wherein a is 0 to 2, C₁₋₄alkoxycarbonyl,

N-(C₁₋₄alkyl)sulphamoyl, *N,N*-(C₁₋₄alkyl)₂sulphamoyl, C₁₋₄alkylsulphonylamino or

(nitrogen-linked heterocyclic ring)carbonyl;

R⁵ and **R⁶** are independently selected from halo, hydroxy, cyano, carbamoyl, ureido,

amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, C₁₋₆alkoxy,

C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, C₁₋₆alkanoylamino, C₂₋₆alkenyloxycarbonyl,

C₁₋₆alkoxycarbonyl, heterocyclyl, heterocycliloxy, heterocyclylamino,

heterocyclyl-*N*-(C₁₋₆alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl, heterocyclylcarbonylamino, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylamino, carbocyclyl-*N*-(C₁₋₆alkyl)amino, carbocyclylsulphonyl, carbocyclylcarbonyl, carbocyclylcarbonylamino, carbocyclyloxycarbonyl, *N*-(C₁₋₆alkyl)amino, 5 *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkoxycarbonylamino, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkylS(O)_a wherein a is 0-2, *N*-(C₁₋₆alkyl)sulphamoylamino, *N,N*-(C₁₋₆alkyl)₂sulphamoylamino, C₁₋₆alkylsulphonylamino, (C₁₋₆alkyl)sulphonyl-*N*-(C₁₋₆alkyl)amino, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl; wherein R⁵ and R⁶ may be optionally substituted on carbon by 10 one or more R⁹; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be independently optionally substituted by R¹⁰;

R⁷ and R⁹ are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, trifluoromethyl, trifluoromethoxy, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, aminosulphamoyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy, C₁₋₄alkoxycarbonyl, 15 C₁₋₄alkoxycarbonylamino, C₂₋₄alkenyloxycarbonyl, C₁₋₄alkanoyl, C₁₋₄alkanoylamino, C₁₋₄alkanoyloxy, *N*-(C₁₋₄alkyl)amino, *N,N*-(C₁₋₄alkyl)₂amino, *N*-(C₁₋₄alkyl)carbamoyl, *N,N*-(C₁₋₄alkyl)₂carbamoyl, C₁₋₄alkylS(O)_a wherein a is 0-2, *N*-(C₁₋₄alkyl)sulphamoylamino, *N,N*-(C₁₋₄alkyl)₂sulphamoylamino, (C₁₋₄alkyl)sulphonylamino, (C₁₋₄alkyl)sulphonyl-*N*-(C₁₋₄alkyl)amino, *N*-(C₁₋₄alkyl)sulphamoyl, 20 *N,N*-(C₁₋₄alkyl)₂sulphamoyl, heterocyclyl, heterocyclyloxy, heterocyclylamino, heterocyclyl-*N*-(C₁₋₄alkyl)amino, heterocyclylsulphonyl, heterocyclylcarbonyl, heterocyclylcarbonylamino, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylamino, carbocyclyl-*N*-(C₁₋₄alkyl)amino, carbocyclylsulphonyl, carbocyclylcarbonyl, carbocyclylcarbonylamino, and carbocyclyloxycarbonyl; wherein R⁷ and 25 R⁹ may be independently optionally substituted on carbon by one or more R¹¹;

R⁸ and R¹⁰ are independently selected from C₁₋₄alkyl, C₁₋₄alkanoyl, sulphamoyl, C₁₋₄alkylsulphonyl, C₁₋₄alkoxycarbonyl, carbamoyl, *N*-(C₁₋₄alkyl)carbamoyl, *N,N*-(C₁₋₄alkyl)₂carbamoyl, *N*-(C₁₋₄alkyl)sulphamoyl, *N,N*-(C₁₋₄alkyl)₂sulphamoyl, heterocyclyl, heterocyclylcarbonyl, heterocyclyloxycarbonyl, heterocyclylsulphonyl, 30 carbocyclyl, carbocyclylcarbonyl, carbocyclyloxycarbonyl and carbocyclylsulphonyl; wherein R⁸ and R¹⁰ may be independently optionally substituted on carbon by one or more R¹²;

- 112 -

R^{11} and R^{12} are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, vinyl, allyl, methoxy, ethoxy, vinyloxy, allyloxy, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, *N*-methylcarbamoyl, *N,N*-dimethylcarbamoyl, methylthio, methylsulphinyl, mesyl, *N*-methylsulphamoyl, *N,N*-dimethylsulphamoyl, heterocyclyl, heterocyclyloxy, heterocyclylcarbonyl, heterocyclyloxycarbonyl, carbocyclyl, carbocyclyloxy, carbocyclylcarbonyl and carbocyclyloxycarbonyl;

x is 0 - 4; wherein the values of R^1 may be the same or different;

y is 0 - 3; wherein the values of R^2 may be the same or different;

10 z is 0 - 3; wherein the values of R^4 may be the same or different; and

n is 0 - 2;

with the providing that when R^3 is a nitrogen linked heterocyclyl, A is a direct bond; or a pharmaceutically acceptable salt, prodrug or solvate thereof.

15 19. A method of treating eating disorders in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB') or a pharmaceutically acceptable salt, prodrug or solvate thereof, as claimed in claim 18.

20 20. A method as claimed in claim 19 wherein the eating disorders are obesity and related disorders, bulimia or anorexia wherein the "related disorders" are diabetes dyslipidaemia, hypertension and sleep disturbances.

21. A method for promoting weight loss in a warm-blooded animal, such as a human
25 being, in need of such treatment which comprises administering to said animal a therapeutically effective amount of a compound of formula (IB') or a pharmaceutically acceptable salt, prodrug or solvate thereof, as claimed in claim 18.

22. The use of a compound of formula (IB') or a pharmaceutically acceptable salt,
30 prodrug or solvate thereof, as claimed in claim 18, in the manufacture of a medicament for the treatment of disorders mediated by the neuropeptide Y5 receptor in a warm-blooded animal, such as a human being.

23. The use of a compound of formula (IB') or a pharmaceutically acceptable salt, prodrug or solvate thereof, as claimed in claim 18, in the manufacture of a medicament for the treatment of eating disorders in a warm-blooded animal, such as a human being.

5

24. A use as claimed in claim 23 wherein the eating disorders are obesity and related disorders, bulimia or anorexia wherein the "related disorders" are diabetes dyslipidaemia, hypertension and sleep disturbances.

10 25. The use of a compound of formula (IB') or a pharmaceutically acceptable salt, prodrug or solvate thereof, as claimed in claim 18, in the manufacture of a medicament for promoting weight loss in a warm-blooded animal, such as a human being.

15 26. A pharmaceutical composition comprising a compound of formula (IB'), or a pharmaceutically acceptable salt, prodrug or solvate thereof, as claimed in claim 18, in association with a pharmaceutically acceptable diluent or carrier for the treatment of disorders mediated by the neuropeptide Y5 receptor in a warm-blooded animal, such as a human being in need of such treatment.

20 27. According to an additional aspect of the invention there is provided a pharmaceutical composition comprising a compound of formula (IB'), or a pharmaceutically acceptable salt, prodrug or solvate thereof, as claimed claim 18, in association with a pharmaceutically acceptable diluent or carrier for the treatment of eating disorders in a warm-blooded animal, such as a human being.

25

28. A pharmaceutical composition comprising a compound of formula (IB'), or a pharmaceutically acceptable salt, prodrug or solvate thereof, as claimed in claim 18, in association with a pharmaceutically acceptable diluent or carrier for promoting weight loss in a warm-blooded animal, such as a human being.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/01899

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D333/76 A61K31/381 A61K31/453 A61K31/454 A61K31/4436
 A61K31/4439 A61P9/00 A61P9/12 A61P3/04 C07D409/12
 C07D413/12 C07D521/00 C07D417/12 C07D409/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, WPI Data, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 882 707 A (ASAHI CHEMICAL IND) 9 December 1998 (1998-12-09) abstract; claims page 49; example 28 ---	1,16-28
A	EP 0 997 458 A (ASAHI CHEMICAL IND) 3 May 2000 (2000-05-03) abstract; claims ---	1,16-28
A	WO 97 20821 A (CIBA GEIGY AG ; RUEEGER HEINRICH (CH); SCHMIDLIN TIBUR (CH); RIGOLL) 12 June 1997 (1997-06-12) abstract; claims page 18 -page 52; examples --- -/--	1,16-28

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
 E earlier document but published on or after the international filing date
 L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 O document referring to an oral disclosure, use, exhibition or other means
 P document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
 & document member of the same patent family

Date of the actual completion of the international search

17 August 2001

Date of mailing of the international search report

24/08/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Paisdor, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/01899

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 64394 A (STAMFORD ANDREW W ;DUGAR SUNDEEP (US); SCHERING CORP (US); WU YUSH) 16 December 1999 (1999-12-16) page 1 page 65 -page 78 -----	16-28

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No
PCT/GB 01/01899

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0882707 A	09-12-1998	AU 715233 B	20-01-2000
		AU 1170897 A	01-08-1997
		NO 983197 A	10-09-1998
		US 6037362 A	14-03-2000
		CA 2242351 A	17-07-1997
		CN 1295064 A	16-05-2001
		CN 1209119 A	24-02-1999
		WO 9725311 A	17-07-1997
		JP 3124242 B	15-01-2001
		JP 9249623 A	22-09-1997
		JP 2000239255 A	05-09-2000
		US 6187809 B	13-02-2001
EP 0997458 A	03-05-2000	AU 735249 B	05-07-2001
		AU 8033498 A	25-01-1999
		DE 997458 T	01-03-2001
		NO 996453 A	03-03-2000
		US 6172099 B	09-01-2001
		ES 2149742 T	16-11-2000
		WO 9901431 A	14-01-1999
WO 9720821 A	12-06-1997	AU 7692696 A	27-06-1997
		ZA 9610021 A	01-06-1997
WO 9964394 A	16-12-1999	AU 4317899 A	30-12-1999
		EP 1086078 A	28-03-2001